



Calhoun: The NPS Institutional Archive
DSpace Repository

Theses and Dissertations

1. Thesis and Dissertation Collection, all items

1990-09

A hierarchical Gamma/Weibull regression model for target detection times

Wang, Chia-Fu

Monterey, California: Naval Postgraduate School

<http://hdl.handle.net/10945/34954>

Copyright is reserved by the copyright owner.

Downloaded from NPS Archive: Calhoun



<http://www.nps.edu/library>

Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

AD-A239 462



2

NAVAL POSTGRADUATE SCHOOL
Monterey, California

DTIC
S D
AUG 19 1991



THESIS

A HIERARCHICAL GAMMA/WEIBULL REGRESSION
MODEL
FOR
TARGET DETECTION TIMES

by

Wang Chia-Fu

September, 1990

Thesis Advisor:

Patricia A. Jacobs

Approved for public release; distribution is unlimited.

91-08092



91 8 16 022

Unclassified

security classification of this page

REPORT DOCUMENTATION PAGE

1a Report Security Classification Unclassified			1b Restrictive Markings		
2a Security Classification Authority			3 Distribution Availability of Report		
2b Declassification Downgrading Schedule			Approved for public release; distribution is unlimited.		
4 Performing Organization Report Number(s)			5 Monitoring Organization Report Number(s)		
6a Name of Performing Organization Naval Postgraduate School		6b Office Symbol (if applicable) 30	7a Name of Monitoring Organization Naval Postgraduate School		
6c Address (city, state, and ZIP code) Monterey, CA 93943-5000		7b Address (city, state, and ZIP code) Monterey, CA 93943-5000			
8a Name of Funding Sponsoring Organization		8b Office Symbol (if applicable)	9 Procurement Instrument Identification Number		
8c Address (city, state, and ZIP code)		10 Source of Funding Numbers			
		Program Element No	Project No	Task No	Work Unit Accession No
11 Title (include security classification) A HIERARCHICAL GAMMA/WEIBULL REGRESSION MODEL FOR TARGET DETECTION TIMES					
12 Personal Author(s) Wang, Chia-Fu					
13a Type of Report Master's Thesis		13b Time Covered From To		14 Date of Report (year, month, day) September 1990	15 Page Count 116
16 Supplementary Notation The views expressed in this thesis are those of the author and do not reflect the official policy or position of the Department of Defense or the U.S. Government.					
17 Cosati Codes			18 Subject Terms (continue on reverse if necessary and identify by block number)		
Field	Group	Subgroup	MAXIMUM LIKELIHOOD ESTIMATE, GAMMA/WEIBULL REGRESSION, NEWTON PROCEDURE, BISECTION SEARCH		
19 Abstract (continue on reverse if necessary and identify by block number)					
<p>Combat models often involve target detection times which may vary with different observers due to characteristics of personnel, or detection systems. They may also be affected by different environmental factors such as visual levels, sea states, terrains, etc. There is often interest in quantifying the effects of different observer characteristics and environmental factors on detection times. A hierarchical gamma/Weibull regression model is considered which can incorporate observer characteristics and environmental effects which may influence the time to detect targets. Numerical procedures for the estimation of parameters of the hierarchical gamma Weibull model based on maximum likelihood are described. Results of simulation experiments to study small sample behavior of the estimates are reported.</p>					
20 Distribution Availability of Abstract			21 Abstract Security Classification		
<input checked="" type="checkbox"/> unclassified unlimited <input type="checkbox"/> same as report <input type="checkbox"/> DTIC users			Unclassified		
22a Name of Responsible Individual Patricia A. Jacobs			22b Telephone (include Area code) (408) 646-2258	22c Office Symbol 62Pw	

Approved for public release; distribution is unlimited.

A Hierarchical Gamma/Weibull Regression Model

For

Target Detection Times

by

Wang, Chia-Fu

Lieutenant , Republic of China Navy

Naval Academy,1986

Submitted in partial fulfillment
of the requirements for the degree of

MASTER OF SCIENCE IN OPERATIONS RESEARCH

from the

NAVAL POSTGRADUATE SCHOOL

September 1990

Author:

Wang, Chia-Fu

Approved by:

Patricia A. Jacobs, Thesis Advisor

D.P. Gaver, Second Reader

Peter Purdue, Chairman
Department of Operations Research

ABSTRACT

Combat models often involve target detection times which may vary with different observers due to characteristics of personnel, or detection systems. They may also be affected by different environmental factors such as visual levels, sea states, terrains, etc. There is often interest in quantifying the effects of different observer characteristics and environmental factors on detection times. A hierarchical gamma/Weibull regression model is considered which can incorporate observer characteristics and environmental effects which may influence the time to detect targets. Numerical procedures for the estimation of parameters of the hierarchical gamma/Weibull model based on maximum likelihood are described. Results of simulation experiments to study small sample behavior of the estimates are reported.



Accession For	
NTIS	CRA&I
DTIC	TAB
Unannounced	
Justification	
By	
Distribution/	
Availability Code	
Dist	Avail and/or Special
A-1	

TABLE OF CONTENTS

I. INTRODUCTION	1
A. BACKGROUND	1
B. TWO RELATED MODELS	1
C. OBJECTIVE AND METHODOLOGY IN THE COMBINED MODEL	2
II. A SINGLE PARAMETER GAMMA HIERARCHICAL MODEL	4
A. MODEL DESCRIPTION	4
B. MAXIMUM LIKELIHOOD ESTIMATION AND THE BISECTION ROOT SEARCH	6
1. The Likelihood Function	6
2. Bisection Root Search for η	8
a. <u>Initial Estimate for η</u>	10
b. <u>Determination of Bounds for the Root</u>	10
c. <u>Bisection Search</u>	11
C. IMPLEMENTATION OF THE SIMULATION AND RESULTS .	11
1. Simulation	11
a. <u>Input to Simulation</u>	13

b. <u>Simulation of Data for ith Observer</u>	13
c. <u>Initial η and Bisection Search</u>	13
2. Results	16
 III. THE HIERARCHICAL GAMMA/WEIBULL REGRESSION MODEL	17
A. MODEL DESCRIPTION	17
B. MAXIMUM LIKELIHOOD ESTIMATES FOR ALL PARAMETERS	17
1. Initial Estimation of a Simple Weibull Regression Model	18
2. Does the Simple Weibull Regression Adequately Account for the Variability of the Data ?	21
3. A Recursive Estimation Procedure for the Parameters of the Full Hierarchical Model	23
a. <u>Bisection Search with Instability Check for Estimating η</u> .	23
b. <u>The Least Squares Regression Procedure to Solve for $\{\beta_k\}$</u>	25
c. <u>Newton Procedure to Solve for $\{\xi_i\}$</u>	28
C. IMPLEMENTATION OF SIMULATION	31
1. Input to Simulation	31
2. Simulation of Data for the i^{th} Observer	32
3. Initial Estimate of β_k 's and ξ_i in the Simple Weibull Regression Model without the Hierarchical Gamma r.v.'s	33

4. Recursive Procedure to Estimate the Parameters of the Full Hierarchical Gamma/Weibull Model	34
D. RESULTS	37
IV. CONCLUSIONS AND RECOMMENDATIONS	42
LIST OF REFERENCES	44
APPENDIX A. STATISTICS AND GRAPHICS FOR THE ESTIMATE OF η IN THE SINGLE PARAMETER GAMMA MODEL	45
APPENDIX B. STATISTICS AND GRAPHICS FOR ALL THE ESTIMATES IN GAMMA/WEIBULL REGRESSION MODEL	56
1. TABLES AND HISTOGRAMS FOR η	56
2. TABLES AND HISTOGRAMS FOR β_0	63
3. TABLES AND HISTOGRAMS FOR β_1	70
4. TABLES AND HISTOGRAMS FOR β_2	77
5. TABLES AND HISTOGRAMS FOR ξ	84
APPENDIX C. SIMULATION PROGRAM FOR THE ESTIMATE OF η IN THE SINGLE PARAMETER GAMMA MODEL	91

APPENDIX D. SIMULATION PROGRAM FOR THE ESTIMATES OF
ALL THE PARAMETERS IN GAMMA/WEIBULL REGRESSION

MODEL	95
INITIAL DISTRIBUTION LIST	100

List of Tables

Table 1.	Mean Bias, Mean Square Error, Std. Error and Uncensored Level for η at $O = 10$ in the Gamma Model	45
Table 2.	Mean Bias, Mean Square Error, Std. Error and Uncensored Level for η at $O = 15$ in the Gamma Model	45
Table 3.	Tendencies of Mean Bias and Mean Square Error for η at $O = 10$ in the Gamma Model	46
Table 4.	Tendencies of Mean Bias and Mean Square Error for η at $O = 15$ in the Gamma Model	51
Table 5.	Mean Bias, Mean Square Error and Std. Error for η at $O = 10$ in the GAM/WEI Regression Model	56
Table 6.	Mean Bias, Mean Square Error and Std. Error for η at $O = 15$ in the GAM/WEI Regression Model	56
Table 7.	Tendencies of Mean Bias and Mean Square Error for η at $O = 10$ in the GAM/WEI Regression Model	57
Table 8.	Tendencies of Mean Bias and Mean Square Error for η at $O = 15$ in the GAM/WEI Regression Model	60
Table 9.	Mean Bias, Mean Square Error and Std. Error for β_0 at $O = 10$ in the GAM/WEI Regression Model	63
Table 10.	Mean Bias, Mean Square Error and Std. Error for β_0 at $O = 15$ in the GAM/WEI Regression Model	63
Table 11.	Tendencies of Mean Bias and Mean Square Error for β_0 at $O = 10$ in the GAM/WEI Regression Model	64
Table 12.	Tendencies of Mean Bias and Mean Square Error for β_0 at $O = 15$ in the GAM/WEI Regression Model	67
Table 13.	Mean Bias, Mean Square Error and Std. Error for β_1 at $O = 10$ in the GAM/WEI Regression Model	70
Table 14.	Mean Bias, Mean Square Error and Std. Error for β_1 at $O = 15$ in the GAM/WEI Regression Model	70
Table 15.	Tendencies of Mean Bias and Mean Square Error for β_1 at $O = 10$ in the GAM/WEI Regression Model	71
Table 16.	Tendencies of Mean Bias and Mean Square Error for β_1 at $O = 15$ in the	

GAM/WEI Regression Model	74
Table 17. Mean Bias, Mean Square Error and Std. Error for β_2 at $O = 10$ in the GAM/WEI Regression Model	77
Table 18. Mean Bias, Mean Square Error and Std. Error for β_2 at $O = 15$ in the GAM/WEI Regression Model	77
Table 19. Tendencies of Mean Bias and Mean Square Error for β_2 at $O = 10$ in the GAM/WEI Regression Model	78
Table 20. Tendencies of Mean Bias and Mean Square Error for β_2 at $O = 15$ in the GAM/WEI Regression Model	81
Table 21. Mean Bias, Mean Square Error and Std. Error for ξ at $O = 10$ in the GAM/WEI Regression Model	84
Table 22. Mean Bias, Mean Square Error and Std. Error for ξ at $O = 15$ in the GAM/WEI Regression Model	84
Table 23. Tendencies of Mean Bias and Mean Square Error for ξ at $O = 10$ in the GAM/WEI Regression Model	85
Table 24. Tendencies of Mean Bias and Mean Square Error for ξ at $O = 15$ in the GAM/WEI Regression Model	88

List of Figures

Figure 1.	The Tendencies of the Bias of η with changes in OBS and TGT at O=10 in the Gamma Model	46
Figure 2.	The Tendencies of the Bias of η with changes in OBS and TGT at O=10 in the Gamma Model	47
Figure 3.	The Tendencies of the Bias of η with changes in OBS and TGT at O=10 in the Gamma Model	47
Figure 4.	Comparison of the Bias of η between Different numbers of Observers with TGT = 15 at O=10 in the Gamma Model	48
Figure 5.	Comparison of the Bias of η between Different numbers of Observers with TGT = 25 at O=10 in the Gamma Model	49
Figure 6.	Comparison of the Bias of η between Different numbers of Observers with TGT = 35 at O=10 in the Gamma Model	50
Figure 7.	The Tendencies of the Bias of η with changes in OBS and TGT at O=15 in the Gamma Model	51
Figure 8.	The Tendencies of the Bias of η with changes in OBS and TGT at O=15 in the Gamma Model	52
Figure 9.	The Tendencies of the Bias of η with changes in OBS and TGT at O=15 in the Gamma Model	52
Figure 10.	Comparison of the Bias of η between Different numbers of Observers with TGT = 15 at O=15 in the Gamma Model	53
Figure 11.	Comparison of the Bias of η between Different numbers of Observers with TGT = 25 at O=15 in the Gamma Model	54
Figure 12.	Comparison of the Bias of η between Different numbers of Observers with TGT = 35 at O=15 in the Gamma Model	55
Figure 13.	Comparison of the Bias of η between Different numbers of Observers with TGT=15 at O=10 in the GAM/WEI Regression Model	57
Figure 14.	Comparison of the Bias of η between Different numbers of Observers with TGT=25 at O=10 in the GAM/WEI Regression Model	58
Figure 15.	Comparison of the Bias of η between Different numbers of Observers with TGT=35 at O=10 in the GAM/WEI Regression Model	59
Figure 16.	Comparison of the Bias of η between Different numbers of Observers	

with TGT = 15 at O = 15 in the GAM/WEI Regression Model	60
Figure 17. Comparison of the Bias of η between Different numbers of Observers with TGT = 25 at O = 15 in the GAM/WEI Regression Model	61
Figure 18. Comparison of the Bias of η between Different numbers of Observers with TGT = 35 at O = 15 in the GAM/WEI Regression Model	62
Figure 19. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 15 at O = 10 in the GAM/WEI Regression Model	64
Figure 20. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 25 at O = 10 in the GAM/WEI Regression Model	65
Figure 21. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 35 at O = 10 in the GAM/WEI Regression Model	66
Figure 22. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 15 at O = 15 in the GAM/WEI Regression Model	67
Figure 23. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 25 at O = 15 in the GAM/WEI Regression Model	68
Figure 24. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 35 at O = 15 in the GAM/WEI Regression Model	69
Figure 25. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 15 at O = 10 in the GAM/WEI Regression Model	71
Figure 26. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 25 at O = 10 in the GAM/WEI Regression Model	72
Figure 27. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 35 at O = 10 in the GAM/WEI Regression Model	73
Figure 28. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 15 at O = 15 in the GAM/WEI Regression Model	74
Figure 29. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 25 at O = 15 in the GAM/WEI Regression Model	75
Figure 30. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 35 at O = 15 in the GAM/WEI Regression Model	76
Figure 31. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 15 at O = 10 in the GAM/WEI Regression Model	78
Figure 32. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 25 at O = 10 in the GAM/WEI Regression Model	79
Figure 33. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 35 at O = 10 in the GAM/WEI Regression Model	80

Figure 34. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 15 at O = 15 in the GAM/WEI Regression Model	81
Figure 35. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 25 at O = 15 in the GAM/WEI Regression Model	82
Figure 36. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 35 at O = 15 in the GAM/WEI Regression Model	83
Figure 37. Comparison of the Bias of ξ between Different numbers of Observers with TGT = 15 at O = 10 in the GAM/WEI Regression Model	85
Figure 38. Comparison of the Bias of ξ between Different numbers of Observers with TGT = 25 at O = 10 in the GAM/WEI Regression Model	86
Figure 39. Comparison of the Bias of ξ between Different numbers of Observers with TGT = 35 at O = 10 in the GAM/WEI Regression Model	87
Figure 40. Comparison of the Bias of ξ between Different numbers of Observers with TGT = 15 at O = 15 in the GAM/WEI Regression Model	88
Figure 41. Comparison of the Bias of ξ between Different numbers of Observers with TGT = 25 at O = 15 in the GAM/WEI Regression Model	89
Figure 42. Comparison of the Bias of ξ between Different numbers of Observers with TGT = 35 at O = 15 in the GAM/WEI Regression Model	90

ACKNOWLEDGEMENTS

To be an international officer student without family here, it is impossible for me to forget all those who have sincerely helped with this thesis in various ways. Especially, during the last few months while my mother struggled with her cancer in Taiwan, their encouragement and concern make the accomplishment of my thesis possible. I would particularly like to thank the following people :

- Professor P. A. Jacobs for her constant guidance and valuable corrections;
- Professor D. P. Gaver for his thorough review and professional assistance;
- My parents for their earnest support and my fiancée, Hui-Fen, for her endless prayer;
- All the brothers and sisters in church for their love in Christ.

I. INTRODUCTION

A. BACKGROUND

Combat models often involve target detection times which may vary with different observers due to characteristics of personnel, or detection systems. They may also be affected by different environmental factors such as visual levels, sea states, terrains, etc. There is often interest in quantifying the effects of different observer characteristics and environmental factors on detection times. In Chapter III a hierarchical gamma/Weibull regression model [Ref. 1] is considered which can incorporate observer characteristics and environmental effects which influence the time to detect targets. Numerical procedures for the estimation of parameters of the hierarchical gamma/Weibull model based on maximum likelihood are described. Results of simulation experiments to study small sample behavior of the estimators are reported. In the remainder of this chapter, two models related to the one considered in this thesis will be described. Numerical procedures for parametric estimation for these models are studied in Bae [Ref. 2]. Experience gained concerning the numerical stability of these procedures will be described.

B. TWO RELATED MODELS

Two parametric models for the distribution of time to detect targets were considered in a thesis written by a Korean officer, Mr. Bae in Sept., 1989 [Ref. 2].

Model 1 : [Ref. 2: pp. 2-14] supposes that there are M observers. Observer i has parameter A_i which characterizes the ability of the observer to detect a target. Observer i is presented with N_i targets. The parameters of all M observers, A_1, A_2, \dots, A_M , are assumed independent with a common two parameter gamma

distribution. Given A_i , the detection times of observer i are conditionally independent, having Weibull distributions with known shape parameter $e^{-\xi_i}$ and known scale parameter μ_{ij} for $j=1,2,\dots,N_i$. Some of the detection times may be censored. Bae [Ref. 2:pp. 2-20] reports on numerical procedures to find the maximum likelihood estimates for the shape and scale parameters of the gamma distribution.

Model 2 : is a Weibull regression model [Ref.2:pp.21-30]. It assumes observer i has explanatory variables $x_{ij1}, x_{ij2}, \dots, x_{ijp}$ relating to his j^{th} target representing factors which influence his time to detection. The detection times for the observers are independent random variables having Weibull distributions. Again, some of the detection times may be censored. The scale parameter of the Weibull distribution for the detection times of the j^{th} target by observer i is of the form $\mu_{ij} = e^{x_{ij}\beta}$ where $x_{ij}\beta = \sum_{k=1}^p x_{ijk}\beta_k$; the shape parameter is of the form $e^{-\xi_i}$. Bae [Ref. 2:pp. 21-36] reports on numerical procedures to find the maximum likelihood estimates of $\{\xi_i\}$ and $\{\beta_k\}$.

C. OBJECTIVE AND METHODOLOGY IN THE COMBINED MODEL

The main effort of this thesis will be to study a full hierarchical gamma/Weibull regression model which is a combination of the two previous models. Since the models considered in Bae [Ref. 2] will be special cases of the one considered in this thesis, it is expected that instabilities found in the numerical procedures to estimate the parameters in the two previous models will appear for the combined model. Two numerical instabilities that result from the numerical procedures of Bae [Ref. 2] are described below.

In Model 1, a modified Newton-Raphson method [Ref. 3] is applied to solve the nonlinear system of equations for the maximum likelihood estimates of the

gamma parameters. It is found that, particularly for small sample sizes, the slope of the likelihood surface is very flat resulting in numerical instability for estimation of the scale parameter η . As a result, in the full hierarchical model presented here we have used a *single parameter gamma distribution* as the second stage model. The maximum likelihood estimate of the parameter is found by the bisection root search method [Ref. 4].

A numerical overflow phenomenon was occasionally found in the Weibull regression model while iterating the Newton procedure for $\{\xi_i\}$ in Model 2 [Ref.2:p.25]. Once again, the flatness of the relationship determining the estimate of $\{\xi_i\}$ resulted in this instability. As a result, the Newton procedure to determine $\{\xi_i\}$ in the gamma/Weibull regression model of this thesis has some checks to detect this numerically instability.

In Chapter II, a numerical procedure is presented to find the maximum likelihood estimate for a *single parameter gamma version* of Model 1. In Chapter III, the full hierarchical model is presented and numerical procedures to estimate the parameters given. In both chapters, simulation results are presented to study the small sample behavior of the estimates. All simulations were carried out on an IBM 3179 G mainframe computer at the Naval Postgraduate School using the APL GRAFSTAT random number package [Ref. 5],[Ref. 6].

We hope that the hierarchical gamma/Weibull regression model and these estimation procedures will be a useful tool to describe and predict target detection times, which are one aspect of the effect of human performance on the battlefield.

II. A SINGLE PARAMETER GAMMA HIERARCHICAL MODEL

A. MODEL DESCRIPTION

This model is very similar to Model 1 in [Ref. 2]¹. The difference is that the parameters A_i , $i = 1, 2, \dots, M$, which reflect the abilities of observer i to detect targets, are assumed to be independent having a *single-parameter gamma distribution*, $GAM(\alpha, \alpha)$ rather than having two parameters. The form of gamma density function used here is

$$g(\theta) = \frac{\alpha(\alpha\theta)^{\alpha-1}}{\Gamma(\alpha)} \cdot e^{-\alpha\theta} \quad \text{for } \theta > 0. \quad (2.1)$$

For numerical reasons we will parametrize α as e^η . For convenience we will reiterate the remainder of the assumptions of Model 1 given in [Ref. 2]. Given $A_i = \theta$, the times it takes for observer i to detect target j , $j = 1, 2, \dots, N_i$, denoted by U_{ij} , are assumed to be conditionally independent random variables with Weibull distributions, $WEI(\mu_{ij}, e^{-\xi_i})$ having cumulative distribution function

$$P\{U_{ij} \leq t | A_i = \theta\} = 1 - \exp\left\{-\theta(t/\mu_{ij})^{e^{-\xi_i}}\right\} \quad t \geq 0 \quad (2.2)$$

independent of other observers. The variations of the A_i are introduced to represent the individual differences between the observers. When the i^{th} observer is presented with his j^{th} target, only an opportunity time O_{ij} is allowed for him to detect it. An observer either successfully detects the target within this time or never detects it.

¹ In the model of [Ref. 2], a two parameter gamma distribution is used.

Data for the i^{th} observer consist of times of detection for the successes and the lengths of opportunity times for the failures. For each $i = 1, 2, \dots, M$, $j = 1, 2, \dots, N_i$, let

$$Y_{ij} = \min(\ln U_{ij}, \ln O_{ij}) \quad (2.3)$$

and

$$\Delta_{ij} = \begin{cases} 1 & \text{if } U_{ij} \leq O_{ij} \\ 0 & \text{otherwise} \end{cases} \quad (2.4)$$

The Y_{ij} are the censored ln-detection times and Δ_{ij} is an indicator of whether the ln-time to detect the j^{th} target by the i^{th} observer is censored or not. Let

$$C_i = \sum_{j=1}^{N_i} \Delta_{ij} \quad (2.5)$$

be the number of targets detected by observer i . In this chapter, we will assume the Weibull parameters $\{\mu_{ij}\}$ and $\{\xi_i\}$ are known constants, and we are only interested in estimating the gamma parameter η . In the next section, the likelihood function for η is given and the bisection method for finding the maximum likelihood estimate of η is described. In the final section, we present results of a simulation study of the behavior of the estimators for small sample sizes.

B. MAXIMUM LIKELIHOOD ESTIMATION AND THE BISECTION ROOT SEARCH

1. The Likelihood Function

Given $A_i = \theta$, the conditional likelihood function [Ref.2:p.3] for observer i using the censored ln-times Y_{ij} is

$$L_i(\mu, \xi | \theta) = \prod_{j=1}^{N_i} \left[\theta e^{(y_{ij} - \ln \mu_{ij}) e^{-\xi_i}} e^{-\xi_i} \right]^{\Delta_{ij}} \exp \left[-\theta e^{(y_{ij} - \ln \mu_{ij}) e^{-\xi_i}} \right]. \quad (2.6)$$

Let

$$S_i = \sum_{j=1}^{N_i} \exp \left\{ (y_{ij} - \ln \mu_{ij}) e^{-\xi_i} \right\} \quad (2.7)$$

and

$$K_i = \exp \left\{ \sum_{j=1}^{N_i} \Delta_{ij} \left[(y_{ij} - \ln \mu_{ij}) e^{-\xi_i} - \xi_i \right] \right\}. \quad (2.8)$$

Equation (2.6) can be rewritten as follows :

$$L_i(\mu, \xi | \theta) = \theta^{C_i} K_i \exp(-\theta S_i). \quad (2.9)$$

From the equations (2.1) and (2.9), it follows that the unconditional likelihood function for observer i is

$$\begin{aligned}
L_i(\mu, \xi, \alpha) &= \int_0^\infty L_i(\mu, \xi | \theta) g(\theta) d\theta \\
&= \int_0^\infty \theta^{C_i} e^{-\theta S_i} \frac{\alpha (\alpha \theta)^{\alpha-1}}{\Gamma(\alpha)} e^{-\alpha \theta} d\theta \\
&= K_i \frac{\alpha^\alpha}{(S_i + \alpha)^{C_i + \alpha}} \prod_{k=0}^{C_i-1} (\alpha + k) .
\end{aligned} \tag{2.10}$$

In equation (2.10), if $C_i = 0$, then $\prod_{k=0}^{C_i-1} (\alpha + k) = 1$. Recall the parametrization $\alpha = e^\eta$. The unconditional ln-likelihood function for observer i can be rewritten as

$$\begin{aligned}
\ln L_i(\mu, \xi, \eta) &= \\
&\ln K_i + e^\eta \eta - (C_i + e^\eta) \ln(S_i + e^\eta) + \sum_{k=0}^{C_i-1} \ln(e^\eta + k)
\end{aligned} \tag{2.11}$$

where if $C_i = 0$, then $\sum_{k=0}^{C_i-1} \ln(e^\eta + k) = 0$. Since the observers are independent, the unconditional ln-likelihood for all M observers is

$$\begin{aligned}
\ln L &= \sum_{i=1}^M \ln L_i = \\
&\sum_{i=1}^M \left\{ \ln K_i + e^\eta \eta - (C_i + e^\eta) \ln(S_i + e^\eta) + \sum_{k=0}^{C_i-1} \ln(e^\eta + k) \right\} .
\end{aligned} \tag{2.12}$$

In this chapter, we will focus attention only on finding the maximum likelihood estimate of η . The derivative of the ln-likelihood with respect to η is

$$\frac{\partial \ln L}{\partial \eta} = f(\eta) = e^{\eta} \sum_{i=1}^M \left\{ \eta + \frac{S_i - C_i}{S_i + e^{\eta}} - \ln(S_i + e^{\eta}) + \sum_{k=0}^{C_i-1} \frac{1}{e^{\eta} + k} \right\}. \quad (2.13)$$

The maximum likelihood estimate for η is the solution of the equation $f(\eta) = 0$:

2. Bisection Root Search for η

Initially, the Newton procedure was used to numerically solve the equation $f(\eta) = 0$. Unfortunately, the procedure frequently either converged to an unreasonable number or encountered problems of numerical instability. Plots of $f(\eta)$ indicate that f can have multiple zeros. A representative plot of f is shown in Figure A² (next page). In Figure A, one ∇ symbol indicates a zero of f , while two ∇ symbols indicate the reasonable root to be used as the estimate for η .

² This is a typical graph for $f(\eta)$ which was generated for simulated data with $O=10$, $M=15$, $N=15$ and random seed = 16807. The true value of η is 1.




Figure A Multiple Roots of the Derivative of the ln-Likelihood Function for η

As a result of these possible multiple zeros, the bisection root search method [Ref.4] is used to solve $f(\eta)=0$. The following sub-section will detail the estimation procedure for η .

a. Initial Estimate for η

The initial estimate for η is similar to its initial estimate in the two-parameter gamma model [Ref. 2], and is obtained by setting ³ $\beta_0 = 0$ in that procedure. Consequentially, the initial estimate of η is

$$\eta^0 = -\ln \left[\hat{m}_2^2 - \hat{m}_1^2 - \frac{\pi^2}{6} \right] \quad (2.14)$$

where

$$\begin{aligned} \hat{m}_1 &= -0.5772 \\ \hat{m}_2 &= \frac{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij} [(y_{ij} - \ln \mu_{ij}) e^{-\xi_{ij}}]^2}{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij}} \end{aligned} \quad (2.15)$$

If $\hat{m}_2^2 - \hat{m}_1^2 - \frac{\pi^2}{6} < 0$, set $\eta^0 = 0$ ⁴.

b. Determination of Bounds for the Root

The purpose of this step is to bound the root that will be used as the estimate between two points, called η_L and η_U . One of these points is η^0 and the other is found as follows :

³ In Mr.Bae's model [Ref2], the regular gamma distribution is parametrized with shape parameter $\nu = e^{\eta + \beta_0}$ and scale parameter $\alpha = e^\eta$. Setting $\beta_0 = 0$, $\alpha = \nu = e^\eta$ yields the one-parameter gamma distribution.

⁴ The estimate $\hat{m}_2^2 - \hat{m}_1^2 - \frac{\pi^2}{6} < 0$ suggests that the variability of the data is small. Hence, it might be better to set η^0 equal to a large positive number, such as 13. A topic for future research is to explore better initial values for η in this default case.

Set $\eta_L = \eta^0 - k \times \delta$
 $\eta_R = \eta^0 + k \times \delta$ $k=1,2,3,\dots$ where δ is a small increment.

Compare the signs of $f(\eta_L)$ and $f(\eta_R)$ with $f(\eta^0)$ until the first different sign from $f(\eta^0)$ is found. This yields a bound on the closest root to η^0 .

c. Bisection Search

After determining the left bound, η_L and right bound, η_R , we begin the bisection search by picking the midpoint between them, called η_M , and computing the value of $f(\eta_M)$. If $f(\eta_M)$ and $f(\eta_L)$ have the same signs, the root is on the other side of η_M ; thus, move the left bound η_L to midpoint η_M . If the signs are different, move the right bound to the midpoint. Iterate the bisection search until $\left| \frac{\eta_L - \eta_R}{\eta_M} \right| < 0.01$. The maximum likelihood estimate of η is η_M .

C. IMPLEMENTATION OF THE SIMULATION AND RESULTS

1. Simulation

In order to study the sampling properties of the maximum likelihood estimate of η , different numbers of observers (M), targets (N_i), and opportunity times (O_{ij}) are used as inputs to the simulation. Each set of inputs (M, N_i, O_{ij}) has a simulation run of (R) 100 replications and each run begins with the same initial random seed (466801743). Statistics, of mean bias (M.B), mean square error (M.S.E) and their standard errors (S.E(M.B) and S.E(M.S.E)), are computed according to following definitions :

$$M.B = \frac{1}{R} \sum_{r=1}^R (\hat{\eta}_r - \eta) \quad (2.16a)$$

$$S.E(M.B) = \sqrt{\frac{1}{R(R-2)} \sum_{r=1}^R ((\hat{\eta}_r - \eta) - M.B)^2} \quad (2.16b)$$

$$M.S.E = \frac{1}{R} \sum_{r=1}^R (\hat{\eta}_r - \eta)^2 \quad (2.17a)$$

$$S.E(M.S.E) = \sqrt{\frac{1}{R(R-2)} \sum_{r=1}^R ((\hat{\eta}_r - \eta)^2 - M.S.E)^2} \quad (2.17b)$$

where $\hat{\eta}_r$ represents the point estimate of parameter η in the r^{th} replication. The fraction of detection level or the averaged uncensoring level (UC) for R (100) replications is determined by

$$UC = \frac{\sum_{r=1}^R \sum_{i=1}^M C_i(r)}{\sum_{r=1}^R \sum_{i=1}^M N_i(r)} \quad (2.18)$$

where $C_i(r)$ is the number of targets detected by observer i in the r^{th} replication and $N_i(r)$ is the number of targets presented to observer i in the r^{th} replication. Theoretically, the longer the opportunity times, the higher the UC level should be. An outline of a replication in the simulation is as follows :

a. Input to Simulation

- M the number of observers (Possible values are 15 , 25 and 35)
- N_i the number of targets presented to the i th observer , where $i = 1, 2, \dots, M$ (Possible values are 15 , 25 and 35)
- O_{ij} the opportunity time for i th observer to detect the j th target, (Possible values are 10 and 15). All of the opportunity times are the same.
- ξ_i the shape parameter of the Weibull distribution for i th observer, where $i = 1, 2, \dots, M$; (Set $\xi_i = 0$ for all i)
- μ_{ij} the scale parameter of the Weibull distribution , $WEI(\mu_{ij}, e^{-\xi_i})$; (Set $\mu_{ij} = 4.2$)
- η the parameter for the single-parameter gamma distribution , $GAM(\alpha, \alpha)$, with $\alpha = e^\eta$; (Set the true value of $\eta = 1$)

b. Simulation of Data for i th Observer

- Generate independent single-parameter gamma r.v.'s A_i from $GAM(\alpha, \alpha)$, having the density function as in equation (2.1).
- Generate independent exponential distribution r.v.'s W_{ij} with mean 1 i.e. $W_{ij} \sim EXP(1)$ for all i and j
- Compute the target detection times : $U_{ij} = \mu_{ij} \left(\frac{W_{ij}}{A_i} \right)^{e^{\xi_i}}$.^s
- Compute the recorded ln-times Y_{ij} as in equation (2.3).
- Compute Δ_{ij} as in equation (2.4).

$$^s \quad P \left\{ \mu_{ij} \left(\frac{W_{ij}}{A_i} \right)^{e^{\xi_i}} \leq t \mid A_i = \theta \right\} = P \left\{ W_{ij} \leq \theta \left(\frac{t}{\mu_{ij}} \right)^{e^{-\xi_i}} \right\} = 1 - \exp \left\{ -\theta \left(\frac{t}{\mu_{ij}} \right)^{e^{-\xi_i}} \right\} \quad \text{which is the}$$

conditional Weibull distribution of equation (2.2).

c. Initial η and Bisection Search

- Compute \hat{m}_2 as in equation (2.15) and determine the initial value of η as in equation (2.14).
- Perform the Bisection root search procedure for the maximum likelihood estimate of η , shown as the flowchart in Figure B.

All these procedures were written in APL codes, named "SIMULA1", "GAMMA", and "FVALUE", which are listed in Appendix C.




Figure B The Algorithm of Bisection Search

2. Results

The results of the simulation experiments appear in Tables and graphs in Appendix A. Tables 1 and 2 contain the general statistics for the bias of $\hat{\eta}$ at the opportunity times $O = 10$ and 15 respectively. Table 3 in combination with box plots, shown in Figures 1 through 3, presents statistics of the bias of the estimates of η as a function of the numbers of targets and observers in the case $O = 10$. Figures 4 to 6 show histograms of the bias with the number of targets fixed and the number of observers changing. Similarly, Table 4 and Figures 7 through 12 focus on the case $O = 15$. In summary, the simulations indicate the following :

- Opportunity time (O)

Tables 1 and 2 indicate that a longer opportunity time is reflected in a higher uncensoring level (UC), which slightly decreases the mean bias (M.B) and mean square error (M.S.E) of η .

- The Number of Observers (M)

Increasing the number of observers for a fixed number of targets results in greater changes in the mean bias and mean square error of η than increasing the number of targets with a fixed number of observers. All the box plots and histograms display this tendency. This phenomenon is quite reasonable, because the parameter η in gamma distribution reflects the variability in abilities of observers. Hence data with more observers will provide better estimation of η (less bias and standard error).

- The Number of Targets (N_i)

The box plots indicate that increasing the number of targets for a fixed number of observers has a small effect on M.B and M.S.E of η with no systematic trend.

III. THE HIERARCHICAL GAMMA/WEIBULL REGRESSION MODEL

A. MODEL DESCRIPTION

Recalling the assumptions of the model, we suppose there are M observers, indexed by i . The i^{th} observer is presented with N_i targets, indexed by j . The variable U_{ij} is the time it takes for observer i to detect the target j . Let $x_{ij1}, x_{ij2}, \dots, x_{ijp}$ be the values of environmental variables (e.g. terrain, sea state, atmospheric condition etc.) which may affect U_{ij} . Given the quantified value of the ability for observer i , $A_i = \theta$, the U_{ij} 's are assumed conditionally independent random variables having Weibull distributions, $WEI(\mu_{ij}, e^{-\tau_i})$, where the scale parameter, instead of being constant, has the form

$$\mu_{ij} = \exp\{x_{ij}\beta\} = \exp\left\{\sum_{k=0}^p x_{ijk}\beta_k\right\}. \quad (3.1)$$

The regression coefficient β_k is the contribution to the Weibull scale parameter of the k^{th} explanatory variable. β_0 is usually the constant term in the regression. The random variables $\{A_i\}$ are assumed independent from a *single parameter gamma distribution*, $GAM(\alpha, \alpha)$ with $\alpha = e^n$.

B. MAXIMUM LIKELIHOOD ESTIMATES FOR ALL PARAMETERS

The model assumptions detailed above are the same as those in Chapter II, except that the scale parameter μ_{ij} in the Weibull distribution is now a function of

the explanatory variables. In this section we consider the estimation of all the parameters of η , $\{\beta_k\}$ and $\{\xi_i\}$.⁶

From equations (2.7), (2.8), (2.12) and equation (3.1), we can rewrite the unconditional ln-likelihood function for all M observers as follows :

$$\ln L = g(\eta, \xi, \beta) = \sum_{i=1}^M \left\{ \ln K_i + e^\eta \eta - (C_i + e^\eta) \ln(S_i + e^\eta) + \sum_{k=0}^{C_i-1} \ln(e^\eta + k) \right\}$$

$$\text{where } K_i = \exp \left\{ \sum_{j=1}^{N_i} \Delta_{ij} \left[(y_{ij} - \mathbf{x}_{ij} \beta) e^{-\xi_i} - \xi_i \right] \right\} \quad (3.2)$$

$$\text{and } S_i = \sum_{j=1}^{N_i} \exp \left\{ (y_{ij} - \mathbf{x}_{ij} \beta) e^{-\xi_i} \right\};$$

Y_{ij} , Δ_{ij} , and C_i are defined in equations (2.3), (2.4) and (2.5) respectively.

In the following sub-sections, we will describe the procedures which are employed to estimate all the model parameters. The general procedure includes the initial rough estimation of $\{\beta_k\}$ and $\{\xi_i\}$ without the hierarchical gamma r.v.'s; a variability check for the need to include the hierarchical gamma parameter η ; and finally, a large recursive procedure for estimating all the model parameters if a hierarchical model is necessary.

1. Initial Estimation of a Simple Weibull Regression Model

An initial model for the data is the simple Weibull regression model for the target detection times described previously as the Model 2 of Bae [Ref. 2:pp. 21-30]. The key steps of the estimation procedure are summarized as follows :

⁶ In the Model 2 of Bae [Ref. 2], no gamma variable was involved. Only the Weibull parameters $\{\beta_k\}$ and $\{\xi_i\}$ were considered.

Step-1.1 : Initially, set ⁷ $\mathbf{x}_{ij}\beta^0 = \ln U_{ij}$ and $\xi_i^0 = 0$.

Step-1.2 : This is an iteratively re-weighted regression step; cf. McCullagh and Nelder [Ref. 7].

Let

$$r_{ij} = (y_{ij} - \mathbf{x}_{ij}\beta^0)e^{-\xi_i^0}; \quad (3.3)$$

$$w_{ij} = \sqrt{\exp\{r_{ij}\}}; \quad (3.4)$$

$$u_{ijh} = x_{ijh}w_{ij}e^{-\xi_i^0}; \quad \text{where } h=0,1,\dots,p \quad (3.5)$$

$$z_{ij} = \frac{(-\Delta_{ij} + w_{ij}^2)}{w_{ij}} + \sum_{h=0}^p u_{ijh}\beta_h^0 = \frac{(-\Delta_{ij} + w_{ij}^2)}{w_{ij}} + w_{ij}e^{-\xi_i^0}\mathbf{x}_{ij}\beta^0. \quad (3.6)$$

Regress the dependent variable z_{ij} against the independent variables u_{ijh} , for $h = 0,1,\dots,p$; e.g. the estimate has the form

$$\underline{B} = (\underline{U}^T \underline{U})^{-1} \underline{U}^T \underline{Z} \quad (3.7)$$

where

⁷ From this page on, all parameters with superscript 0 imply the current values of their estimates.

$$\begin{aligned}
 \mathbf{Z} = & \begin{bmatrix} z_{11} \\ z_{12} \\ \vdots \\ z_{1N_1} \\ z_{21} \\ \vdots \\ z_{2N_2} \\ \vdots \\ z_{M1} \\ \vdots \\ z_{MN_M} \end{bmatrix} \quad \mathbf{U} = \begin{bmatrix} u_{110} & \cdots & u_{11p} \\ u_{120} & \cdots & u_{12p} \\ \vdots & \vdots & \vdots \\ u_{1N_10} & \cdots & u_{1N_1p} \\ u_{210} & \cdots & u_{21p} \\ \vdots & \vdots & \vdots \\ u_{2N_20} & \cdots & u_{2N_2p} \\ \vdots & \vdots & \vdots \\ u_{M10} & \cdots & u_{M1p} \\ \vdots & \vdots & \vdots \\ u_{MN_M0} & \cdots & u_{MN_Mp} \end{bmatrix} \quad (3.8), (3.9)
 \end{aligned}$$

Step-1.3 : Compute the new ξ_i by an approximation based on Newton's procedure. First, update the value of

$$r_{ij} = (y_{ij} - x_{ij} \beta^0) e^{-\xi_i^0}, \quad \text{where } \beta^0 = \underline{B}. \quad (3.10)$$

Compute

$$\xi_i = \xi_i^0 + \frac{\left\{ C_i - \sum_{j=1}^{N_i} r_{ij} [-\Delta_{ij} + \exp(r_{ij})] \right\}}{\left\{ -C_i - \sum_{j=1}^{N_i} r_{ij}^2 \exp(r_{ij}) \right\}} \quad (3.11)$$

where $C_i = \sum_{j=1}^{N_i} \Delta_{ij}$.

Step-1.4 : Update ξ_i and β_k

$$\begin{aligned} \xi_i^0 &= \xi_i \\ \beta_k^0 &= \beta_k \quad \text{where } k=0,1,\dots,p \end{aligned}$$

Compute the new value of $x_{ij}\beta^0 = \sum_{k=0}^p x_{ijk}\beta_k^0$ and iterate the regression, starting at Step-1.2, for one more time.

2. Does the Simple Weibull Regression Adequately Account for the Variability of the Data ?

The full hierarchical model will be used if the data indicate more variability than can be described by the simple Weibull regression. If the data do come from the simple Weibull model, then

$$\left(\frac{U_{ij}}{\mu_{ij}} \right)^{e^{-t_i}} \quad j=1,2,\dots,N_i$$

will have the same distribution as independent exponential random variables⁸ with mean 1. To assess the variability not accounted for by the simple regression model, we compute

$$\hat{\sigma}_E^2 = \frac{1}{(N-1)} \sum_{ij} \left[\left(\frac{U_{ij}}{\mu_{ij}} \right)^{e^{-t_{ij}}} - 1 \right]^2 \quad (3.12)$$

$$\text{and} \quad N = \sum_{i=1}^M N_i .$$

Since the variance of a unit exponential is 1, if $\hat{\sigma}_E^2 \leq 1$, then we will stop the model fitting with the simple Weibull regression model. If $\hat{\sigma}_E^2 > 1$, we go on to the hierarchical model to explain the extra-variability. Unfortunately, crude estimate of $\hat{\sigma}_E^2$ uses both the observed and unobserved U_{ij} . A topic for future research is to improve the estimate of the variability not accounted for by the simple regression model.

Step-1.5 : If $\hat{\sigma}_E^2 \leq 1$, stop and use the Weibull regression model only. Otherwise, the data will be modeled with the full hierarchical model. A recursive procedure to estimate η , $\{\beta_k\}$ and $\{\xi_i\}$ follows and is initialized with $\{\beta_k^0\}$ and $\{\xi_i^0\}$ computed above.

⁸ $P\{U_{ij} \leq t\} = P\{\mu_{ij}(W_{ij})^{e^{-t_{ij}}} \leq t\} = P\{W_{ij} \leq (t/\mu_{ij})^{e^{-t_{ij}}}\} = 1 - e^{-(t/\mu_{ij})^{e^{-t_{ij}}}}$ which is the pure Weibull distribution.

3. A Recursive Estimation Procedure for the Parameters of the Full Hierarchical Model

a. Bisection Search with Instability Check for Estimating η

The partial derivative of equation (3.2) with respect to η , $\frac{\partial \ln L}{\partial \eta}$, is the same as the derivative of the ln-likelihood with respect to η for the model of Chapter II. The derivative appears in (2.13) and once again the problem is to solve the equation $f(\eta)=0$. Using the estimates $\{\beta_k^0\}$ and $\{\xi_i^0\}$ an initial estimate for η is computed as in equation (2.14) ; that is ,

$$\eta^0 = -\ln \left[\hat{m}_2^2 - \hat{m}_1^2 - \frac{\pi^2}{6} \right] \quad (3.13)$$

where

$$\begin{aligned} \hat{m}_1 &= -0.5772 \\ \hat{m}_2 &= \frac{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij} \left[(y_{ij} - x_{ij} \beta^0) e^{-x_{ij}^2} \right]^2}{\sum_{i=1}^M \sum_{j=1}^{N_i} \Delta_{ij}}. \end{aligned} \quad (3.14)$$

If $\hat{m}_2^2 - \hat{m}_1^2 - \frac{\pi^2}{6} < 0$, set $\eta^0 = 0$.⁹

In addition to determining the bounds for a reasonable root to be the estimate of η as described in Chapter II, a check for the shape of $f(\eta)$ is also performed. This additional check is performed because two of the simulation replications exhibited a very flat $f(\eta)$. The graph of $f(\eta)$ for one of the two

⁹ Same as footnote 4 on page 10.

replications appears in Figure C¹⁰. This unusual shape for $f(\eta)$ results in the bisection method converging to an unreasonable root (e.g. $\hat{\eta} = 146$ in the Figure C).




Figure C An Ill-Conditioned Derivative of the ln-Likelihood Function for η

The graph in Figure C indicates that the values of $f(\eta)$ are asymptotic to a fixed nonzero value as η tends to $-\infty$; this behavior is the same as in Figure A. However, when η is positive, f has very flat curve. The values of $f(\eta)$ are positive

¹⁰ The graph is one of the only two ill-conditioned cases for $f(\eta)$ in the simulations. This case occurred in the gamma/Weibull regression model with $O=10$, $M=15$, $N=25$ and random seed = 1905665785. The other case occurred with $O=15$, $M=15$, $N=25$ and same random seed. The true value of η is 1.

and close to zero but the curve does not actually cross zero until η is large. The following check is done to detect this situation. If a reasonable right hand bound for a zero of $f(\eta)$ cannot be found, check the values of $f(\eta)$ and $f'(\eta)$ for η positive. If the absolute value of $f(\eta)$ is less than ϵ_1 and the absolute value of $f'(\eta)$ is less than ϵ_2 , where $1 \gg \epsilon_1 > \epsilon_2 > 0$, then the simulation stops for this set of random numbers and new random numbers are generated. Otherwise, perform the bisection search as described in Chapter II. In the procedure, $f'(\eta)$ for η positive is estimated as

$$f'(\eta_{R(k)}) \approx \frac{f(\eta_{R(k)}) - f(\eta_{R(k-1)})}{\delta} \quad (3.15)$$

$$\text{where } \eta_{R(k)} = \eta^0 + k \times \delta \\ \eta_{R(k-1)} = \eta^0 + (k-1) \times \delta.$$

The algorithm of this bisection search with the check for flatness of the curve of f will be detailed later in Figure D.

b. The Least Squares Regression Procedure to Solve for $\{\beta_k\}$

The previous estimates of η^0 , $\{\beta_k^0\}$ and $\{\xi_i^0\}$ are input to this stage.

The partial derivative of equation (3.2) with respect to β_k is

$$\frac{\partial \ln L}{\partial \beta_k} = \sum_{i=1}^M \left\{ \frac{\partial \ln K_i}{\partial \beta_k} + \frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \cdot \frac{\partial S_i}{\partial \beta_k} \right\} \quad (3.16)$$

where

$$\frac{\partial \ln K_i}{\partial \beta_k} = \sum_{j=1}^{N_i} \Delta_{ij} (-x_{ijk}) e^{-\xi_i^0}$$

$$\frac{\partial S_i}{\partial \beta_k} = \sum_{j=1}^{N_i} \exp[(y_{ij} - x_{ij} \beta) e^{-\xi_i^0}] (-x_{ijk} e^{-\xi_i^0}) .$$

Thus

$$\frac{\partial \ln L}{\partial \beta_k} = \sum_{i=1}^M \sum_{j=1}^{N_i} \left\{ -\Delta_{ij} + \frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \exp[(y_{ij} - x_{ij} \beta) e^{-\xi_i^0}] \right\} (x_{ijk} e^{-\xi_i^0}) . \quad (3.17)$$

Two terms of a Taylor expansion of equation (3.17) yields

$$\begin{aligned} 0 = \frac{\partial \ln L}{\partial \beta_k} &\sim \sum_{i=1}^M \sum_{j=1}^{N_i} \left\{ \left[-\Delta_{ij} + \frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \exp[(y_{ij} - x_{ij} \beta^0) e^{-\xi_i^0}] \right] \right. \\ &\quad \left. - \left\{ \exp[(y_{ij} - x_{ij} \beta^0) e^{-\xi_i^0}] \sum_{h=0}^P x_{ijk} e^{-\xi_i^0} (\beta_h - \beta_h^0) \right\} \right\} (x_{ijk} e^{-\xi_i^0}) . \end{aligned} \quad (3.18)$$

Let

$$w_{ij} = \sqrt{\frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \exp[(y_{ij} - x_{ij} \beta^0) e^{-\xi_i^0}]} . \quad (3.19)$$

The equation (3.18) can be rewritten as

$$0 = \frac{\partial \ln L}{\partial \beta_k} = \sum_{i=1}^M \sum_{j=1}^{N_i} \left(\frac{-\Delta_{ij} + w_{ij}^2}{w_{ij}} - \sum_{h=0}^p x_{ijh} e^{-\xi_i^0} (\beta_h - \beta_h^0) \right) (x_{ijk} e^{-\xi_i^0}) \quad (3.20)$$

Let

$$u_{ijh} = x_{ijh} w_{ij} e^{-\xi_i^0} \quad (3.21)$$

and

$$z_{ij} = \frac{(-\Delta_{ij} + w_{ij}^2)}{w_{ij}} + \sum_{h=0}^p u_{ijh} \beta_h^0 = \frac{(-\Delta_{ij} + w_{ij}^2)}{w_{ij}} + w_{ij} e^{-\xi_i^0} x_{ij} \beta^0 \quad (3.22)$$

The equation (3.20) can be rewritten as

$$0 = \sum_{i=1}^M \sum_{j=1}^{N_i} \left(z_{ij} - \sum_{h=0}^p u_{ijh} \beta_h \right) u_{ijk} \quad (3.23)$$

which are the normal equations for a least squares regression having dependent variable z_{ij} and independent variables u_{ijk} , $k=0,1,\dots,p$.

The solution of (3.23) is

$$\underline{B} = (\underline{U}^T \underline{U})^{-1} \underline{U}^T \underline{Z} \quad (3.24)$$

where \underline{Z} and \underline{U} are defined as in equations (3.25) and (3.26) respectively.

$$\begin{aligned}
 \mathbf{Z} = \begin{bmatrix} z_{11} \\ z_{12} \\ \vdots \\ z_{1N_1} \\ z_{21} \\ \vdots \\ z_{2N_2} \\ \vdots \\ z_{M1} \\ \vdots \\ z_{MN_M} \end{bmatrix} \quad \mathbf{U} = \begin{bmatrix} u_{110} & \cdots & u_{11p} \\ u_{120} & \cdots & u_{12p} \\ \vdots & \vdots & \vdots \\ u_{1N_1 0} & \cdots & u_{1N_1 p} \\ u_{210} & \cdots & u_{21p} \\ \vdots & \vdots & \vdots \\ u_{2N_2 0} & \cdots & u_{2N_2 p} \\ \vdots & \vdots & \vdots \\ u_{M10} & \cdots & u_{M1p} \\ \vdots & \vdots & \vdots \\ u_{MN_M 0} & \cdots & u_{MN_M p} \end{bmatrix} \quad (3.25), (3.26)
 \end{aligned}$$

This procedure is an iteratively re-weighted least squares regression; cf. McCullagh and Nelder [Ref. 7].

c. Newton Procedure to Solve for $\{\xi_i\}$

The prior estimates of η^0 , $\{\beta_k^0\}$, $\{\xi_i^0\}$ and $\{\xi_i^0\}$ are input to this stage. In order to update the values of $\{\xi_i\}$, we need to have the partial derivative of unconditional ln-likelihood function for observer i with respect to ξ_i . The unconditional ln-likelihood function for observer i is the same as equation (2.11).

The derivative is as follows :

$$\frac{\partial \ln L_i}{\partial \xi_i} = h(\xi_i) = \left\{ \sum_{j=1}^{N_i} \Delta_{ij} \left[(y_{ij} - x_{ij} \beta^0) (e^{-\xi_i}) - 1 \right] \right\} - \frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \cdot \frac{\partial S_i}{\partial \xi_i} \quad (3.27)$$

where $\frac{\partial S_i}{\partial \xi_i} = \sum_{j=1}^{N_i} \exp \left\{ (y_{ij} - x_{ij} \beta^0) e^{-\xi_i} \right\} (-e^{-\xi_i}) (y_{ij} - x_{ij} \beta^0) .$

Again, let

$$r_{ij} = (y_{ij} - x_{ij} \beta^0) e^{-\xi_i} .$$

The equation (3.27) can be rewritten as

$$\frac{\partial \ln L_i}{\partial \xi_i} = h(\xi_i) = -C_i + \sum_{j=1}^{N_i} r_{ij} \left[-\Delta_{ij} + \frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \exp(r_{ij}) \right] . \quad (3.28)$$

Treating $\frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}}$ as a constant, then

$$\frac{\partial^2 \ln L_i}{\partial \xi_i^2} = \frac{\partial h(\xi_i)}{\partial \xi_i} = -C_i - \frac{C_i + e^{\eta^0}}{S_i + e^{\eta^0}} \sum_{j=1}^{N_i} \exp(r_{ij}) r_{ij}^2 \quad (3.29)$$

If $\frac{\partial \ln L_i}{\partial \xi_i} = h(\xi_i) = 0$, the Newton equation for ξ_i is

$$0 = \frac{\partial \ln L_i}{\partial \xi_i} = h(\xi_i) = h(\xi_i^0) + \frac{\partial}{\partial \xi_i} h(\xi_i^0) [\xi_i - \xi_i^0] . \quad (3.30)$$

Update the value of ξ_i by solving the following equation :

$$\xi_i = \xi_i^0 - \frac{h(\xi_i^0)}{\frac{\partial}{\partial \xi_i} h(\xi_i^0)} . \quad (3.31)$$

As noted before, it is possible for equation (3.31) to be badly conditioned. If $|h(\xi_i^0)| < \epsilon_1$ and $\left| \frac{\partial}{\partial \xi_i} h(\xi_i^0) \right| < \epsilon_2$, where $1 > \epsilon_1 > \epsilon_2 > 0$, then equation (3.31) is badly conditioned. This behavior happens occasionally and occurs for only one ξ_i out of the M . To avoid this behavior, if $|h(\xi_i^0)| < \epsilon_1$ and $\left| \frac{\partial}{\partial \xi_i} h(\xi_i^0) \right| < \epsilon_2$, then put $\xi_i = \xi_i^0$; otherwise let ξ_i be the solution to equation (3.31). Return to the bisection search procedure and iterate steps 3.a to 3.c until

$$\max \left(\left| \frac{\eta_L - \eta_R}{\eta_M} \right|, \left| \frac{\beta_k - \beta_k^0}{\beta_k^0} \right|, \left| \frac{\xi_i - \xi_i^0}{\xi_i^0} \right| \right) < 0.01 \text{ for all } k=0, \dots, p \text{ and } i = 1, \dots, M.$$

C. IMPLEMENTATION OF SIMULATION

Simulation will be used to study the small sample behavior of the estimators of the five parameters of the gamma/Weibull model. Different numbers of observers (M), targets (N_i) and opportunity times (O_{ij}) are used as the input to the simulations. For each set of input, (O, M, N_i), the simulation run starts with the same random seed as in Chapter II. The simulation is run until there are 100 (R) estimates of η . A run will contain more than 100 replications if the simple Weibull regression model is found to describe the variability of the data for some of the replications or if the procedure to find the estimate of η is badly conditioned. The statistics of mean bias (M.B), mean square error (M.S.E) and their respective standard errors (S.E(M.B) and S.E(M.S.E)) for all the estimates are computed using the definitions in equations (2.16a,b) and (2.17a,b). The averaged uncensoring levels (UC) as computed using equation (2.18) are also reported. The statistics for $\{\xi_i\}$ do not include those ξ_i for which $|h(\xi_i)| < \epsilon_1$, ($\epsilon_1=0.1$), and $|h'(\xi_i)| < \epsilon_2$, ($\epsilon_2=0.01$).

The remainder of this section provides an outline of one replication in the simulation.

1. Input to Simulation

- M the number of observers (Possible values are 15 , 25 and 35).
- ξ_i the shape parameter of the Weibull distribution for i th observer, where $i = 1, 2, \dots, M$; (Set $\xi_i = 0$ for all i).
- η the parameter for the single-parameter gamma distribution, $GAM(\alpha, \alpha)$, where $\alpha = e^\eta$; (Set $\eta = 1$).
- N_i the number of targets presented to the i th observer , where $i = 1, 2, \dots, M$; (Possible values are 15 , 25 and 35).

- x_{ijk} the k^{th} environmental factor which affects the observation time of the i^{th} observer when presented with the j^{th} target.
- O_{ij} the opportunity time for the i^{th} observer to detect the j^{th} target; (Possible values are 10 and 15).
- $\{\beta_k\}$ regression coefficients for the scale parameter of Weibull distribution, $WEI(\mu_{ij}, e^{-t_i})$, where $k = 0, 1, 2, \dots, p$ and

$$\mu_{ij} = \exp\left\{\sum_{k=0}^p x_{ijk} \beta_k\right\}; \text{ (set } p = 2; \beta_0 = 0.8, \beta_1 = -0.2 \text{ and } \beta_2 = 0.5 \text{).}$$

2. Simulation of Data for the i^{th} Observer

- Generate a single-parameter gamma r.v. A_i from $GAM(\alpha, \alpha)$, having density function as in equation (2.1) where $\alpha = e^\eta$ and $\eta = 1$.
- Generate environmental factors r.v. x_{ijk} ; the x_{ijk} are independent with normal distributions having mean μ_k and variance σ_k^2 for $k = 1, 2, \dots, p$; (Set $p = 2$; $\mu_1 = 1$, $\sigma_1^2 = 0.5$ and $\mu_2 = 2$, $\sigma_2^2 = 1$). Put $x_{ij0} = 1$ for all i, j .
- Compute the Weibull shape parameter μ_{ij} as in equation (3.1).
- Generate independent exponential distribution r.v.'s W_{ij} with mean 1; i.e. $W_{ij} \sim EXP(1)$ for all i and j .
- Compute the target detection times: $U_{ij} = \mu_{ij} \left(\frac{W_{ij}}{A_i} \right)^{t_i}$.
- Compute the recorded ln-times Y_{ij} , censoring indicator Δ_{ij} and C_i using the definitions in equation (2.3), (2.4) and (2.5) respectively.

3. Initial Estimate of β_k 's and ξ_i in the Simple Weibull Regression Model without the Hierarchical Gamma r.v.'s

Step-1.1 : Initially, set $x_{ij}\beta^0 = \ln U_{ij}$ and $\xi_i^0 = 0$

Step-1.2 : Iteration for regression :

- Compute r_{ij} , w_{ij} , u_{ij} and z_{ij} as in equations (3.3) to (3.6).
- Regress the dependent variable z_{ij} against the independent variables u_{ijh} for $\{\beta_k\}$ as in equations (3.7) to (3.9).

Step-1.3 : Newton's procedure for estimating ξ_i

- Update the value of r_{ij} as in equation (3.10).
- Compute the new ξ_i as in equation (3.11) with no need to check the ill condition.

Step-1.4 Put $\{\xi_i^0\} = \{\xi_i\}$ and $\{\beta_k^0\} = \{\beta_k\}$ for all $i=1,2,\dots,M$ and

$k=0,1,\dots,p$. Compute new value of $x_{ij}\beta^0 = \sum_{k=0}^p x_{ijk}\beta_k^0$ and return to Step-1.2 for only one iteration.

Step-1.5 Check for extra-variability in the data by computing the sample variance $\hat{\sigma}_E^2$ for the times U_{ij} as in equation (3.12).

If $\hat{\sigma}_E^2 \leq 1$, stop and quote the simple Weibull regression model only with the parameters as estimated.

Otherwise, execute next the recursive procedure.

4. Recursive Procedure to Estimate the Parameters of the Full Hierarchical Gamma/Weibull Model

Input : initial β_k^0 , $k=0,1,\dots,p$; ξ_i^0 , $i=1,2,\dots,M$.

Step-2.1 : Initial condition for gamma parameter η

- Update $x_{ij}\beta^0 = \sum_{k=0}^p x_{ijk}\beta_k^0$.
- Find the initial condition for η using equations (3.13) and (3.14)

If $\hat{m}_2^2 - \hat{m}_1^2 - \frac{\pi^2}{6} < 0$, set $\eta^0 = 0$.

Step-2.2 Perform the bisection search for η using the algorithm shown in Figure D (next page). The algorithm includes a check for the flatness of the curve of $f(\eta)$.




Figure D The Algorithm of Bisection Search with Instability Check for η

¹¹ In the Algorithm, we let $f'(\eta_R) \sim \frac{[f(\eta_{R(k)}) - f(\eta_{R(k-1)})]}{\delta}$ where $\eta_{R(k)} = \eta^0 + k \times \delta$
 $\eta_{R(k-1)} = \eta^0 + (k-1) \times \delta$,

and k is the current index of increments.

Step-3.1 : Iteration for the regression coefficients $\{\beta_k\}$

- Compute w_{ij} , u_{ij} and z_{ij} using equations (3.19), (3.21) and (3.22).
- Regress the dependent variable z_{ij} against the independent variables u_{ijh} using equations (3.24), (3.25) and (3.26).

Step-4.1 : Iteration for the ln-shape parameters $\{\xi_i\}$, of the Weibull distribution

- Update $x_{ij}\beta^0 = \sum_{k=0}^p x_{ijk}\beta_k^0$ and $r_{ij} = (y_{ij} - x_{ij}\beta^0)e^{-\xi_i^0}$.
- Compute $h(\xi_i^0)$ and $\frac{\partial}{\partial \xi_i} h(\xi_i^0)$ using equations (3.28) and (3.29) respectively.
- If $|h(\xi_i^0)| < \epsilon_1$ and $\left| \frac{\partial}{\partial \xi_i} h(\xi_i^0) \right| < \epsilon_2$, for $\epsilon_1 = 0.1$ and $\epsilon_2 = 0.01$, keep $\xi_i = \xi_i^0$. Otherwise, update ξ_i using equation (3.31).

Step-4.2 : Test for stop criteria

If $\max \left(\left| \frac{\eta_L - \eta_R}{\eta_M} \right|, \left| \frac{\beta_k - \beta_k^0}{\beta_k^0} \right|, \left| \frac{\xi_i - \xi_i^0}{\xi_i^0} \right| \right) < 0.01$ for all $k=0, \dots, p$ and $i=1, \dots, M$, then, stop and record the estimates. Otherwise, set $\eta^0 = \eta$, $\beta_k^0 = \beta_k$, $\xi_i^0 = \xi_i$ and return to Step-2.2.

The simulation programs for this gamma/Weibull regression model, named "SIMULA2", "GAMWEI" and "FVALUE", are detailed in Appendix D.

D. RESULTS

Results of the simulation experiments are presented in Appendix B. The results are presented in five sections. Each section shows results for one estimator; the gamma parameter η and the Weibull parameters β_0 , β_1 , β_2 and ξ_i . Each section contains tables of statistics and histograms of estimate bias for different numbers of observers (M or OBS), targets (N or TGT) and opportunity times (O).

The first two tables in each section present the statistics for different opportunity times (O=10 and O=15 in our case). The next two tables organize the information to highlight relations between the number of targets and the number of observers and the statistics of the estimates. Histograms of the biases of the estimates are also presented. All parameter estimates appearing in Appendix B are from replications for which the full hierarchical gamma/Weibull is estimated. Since $\xi_i \neq 1$, the statistics for $\{\xi_i\}$ are for the average of the $\{\xi_i\}$ for all the observers in a replication. The average includes only those ξ_i 's for which the Newton procedure converged. The replications for which $\hat{\sigma}_E^2 \leq 1$ and the replications in which η could not be found are not used to compute the statistics appearing in Appendix B; statistics concerning the numbers of these cases appear in Tables A and B below.

The statistics in Appendix B use 100 replications. Tables A and B below show the averaged uncensoring levels, the number of additional replications for which η could not be found due to the flatness of f , and the number of times a ξ_i could not be found using Newton procedure due to the smallness of $|h(\xi_i)|$ and $|h'(\xi_i)|$. The results of Tables A and B (next page) are summarized as follows :

- **Uncensoring level (UC) :**
A longer opportunity time (O) is reflected in a higher average uncensoring level (UC). The opportunity time $O = 10$ gives about 76% UC, while $O = 15$ gives about 85% UC. This behavior is what we would expect.
- **The variability of the data explained by the simple Weibull regression :**
Different parameter estimates in the simple Weibull regression cause the different number of cases for which $\{\hat{\sigma}_E^2 \leq 1\}$ for the same input of number of observers and number of targets in Tables A and B. In both Tables A and B, the cases of $\{\hat{\sigma}_E^2 \leq 1\}$ appear more frequently when the number of observers (M) is smaller. This is reasonable behavior since more observers tend to provide more evidence that a hierarchical model is needed to explain the variability of the data. The manner in which $\hat{\sigma}_E^2$ is computed prevents drawing conclusions concerning the effect of censoring on the assessment of unexplained variability.
- **Instability check for η :**
There are only two replications in which η could not be found using the bisection search for the root of $f(\eta)$. The function $f(\eta)$ for one of them is plotted as in Figure C, and the function for the other is similar. Both replications used the same random seed but had a different set of inputs, (M, N_i, O_{ij}) . These two replications were stopped and new random numbers were drawn. This behavior did not occur in the model of Chapter II with known parameters for the Weibull distribution.
- **Instability check for ξ :**
Smaller opportunity times tend to have more replications in which the procedure to find ξ_i is badly conditioned. The maximum number for any case is 2.

TABLE A : Comparisons of Uncensoring levels, Extra Variability Checks and Checks of Numerical Instability at O = 10

Number of Observers	Number of Targets	Average UC	Number of $\{\hat{\sigma}_E^2 < 1\}$	Num. unstable $f(\eta)$	Num. unstable h and h' ¹²
15	15	76.4%	11	0	1 _a
	25	76.4%	5	1	0
	35	75.6%	6	0	0
25	15	75.8%	1	0	1 _b
	25	75.8%	2	0	1 _c
	35	76.4%	1	0	0
35	15	76.2%	0	0	2 _d
	25	76.3%	1	0	1 _e
	35	76.3%	0	0	0

TABLE B : Comparisons of Uncensoring levels, Extra Variability Checks and Checks of Numerical Instability at O = 15

Number of Observers	Number of Targets	Average UC	Number of $\{\hat{\sigma}_E^2 < 1\}$	Num. unstable $f(\eta)$	Num. unstable h and h'
15	15	85.0%	23	0	0
	25	85.4%	6	1	0
	35	84.6%	11	0	0
25	15	84.5%	3	0	1 _f
	25	84.7%	7	0	0
	35	85.3%	5	0	0
35	15	85.2%	5	0	0
	25	84.9%	3	0	0
	35	85.1%	2	0	0

¹² All the cases of numerically unstable h and h' were obtained while replicating the simulations starting with random seeds a:999445582, b:444977940, c:538790986, d:999445582 & 153499980, e:1011786849 and f:444977940.

In Appendix B, the statistics for the average observer ξ_i do not include those individual ξ_i for which the Newton procedure did not converge. The statistics for β_k do not include those replications for which the simple Weibull regression model was used to describe the data. A summary of the results of the simulation experiment shown in Appendix B follows :

- The estimate of the gamma parameter, $\hat{\eta}$

The mean bias tends to be positive. The histograms of biases indicate that the distribution is skewed to the right. Since the variance of A_i is $e^{-\eta}$, the positive mean bias of η suggests that for small sample sizes the procedure is indicating less between variability than there is in the data. Increasing the number of observers tends to decrease the mean bias and mean square error. Increasing the opportunity time has little effect on mean bias and mean square error. Increasing the number of targets has little effect on mean bias and mean square error. As noted in Chapter II, this behavior is reasonable.

- The estimates of the Weibull parameters, $\hat{\beta}_0$ through $\hat{\beta}_2$

The first two tables in parts 2 through 4 of Appendix B indicate that all the M.B and M.S.E of $\{\beta_k\}$ are quite small compared to those for η . This is due to the larger sample size used in computing $\{\beta_k\}$. An increase in the opportunity time has little effect on the mean bias and mean square error. Increasing the number of targets or observers with a fixed number of the other yields small changes in M.B and M.S.E. However, increasing both the number of targets and the number of observers does tend to decrease the mean bias and mean square error of $\{\beta_k\}$. The M.B's are both positive and negative.

- The average estimate of the Weibull parameter, $\hat{\xi} = \frac{1}{M} \sum_{i=1}^M \hat{\xi}_i$

Since $\xi_i = 1$ in the simulations, the histograms and statistics include the averaged $\hat{\xi}_i$ for all the observers for which the Newton procedure for ξ_i converged. Generally, all the magnitudes of M.B and M.S.E of $\hat{\xi}$ in the first 2 tables of part 5 in Appendix B are quite small. All the M.B of the averaged $\hat{\xi}$ are negative. The M.B. and M.S.E. are slightly smaller for

longer opportunity times. Increasing the number of targets or the number of observers with the other fixed results small changes in the M.B and M.S.E. Increasing both the number of targets and the number of observers tends to decrease the mean bias and mean square error.

IV. CONCLUSIONS AND RECOMMENDATIONS

The main effort of this thesis is to study the small sample behavior of estimators for a hierarchical gamma/Weibull regression model for target detection times. This model can also be used to describe and predict times to failure of similar machines, (e.g. engines) in different environments. In this thesis, the model assumes that there are M observers. The i^{th} observer is presented with N_i targets. The variable U_{ij} is the time it takes for observer i to detect the target j . Let $x_{ij1}, x_{ij2}, \dots, x_{ijp}$ be the values of environmental variables which may affect U_{ij} . Given the quantified value of the ability for observer i , $A_i = \theta$, the U_{ij} 's are assumed conditionally independent random variables having Weibull distributions, $WEI(\mu_{ij}, e^{-\xi_i})$, where the scale parameter has the form $\mu_{ij} = \exp\{x_{ij}\beta\} = \exp\left\{\sum_{k=0}^p x_{ijk}\beta_k\right\}$. The random variables $\{A_i\}$ are assumed independent from a single parameter gamma distribution, $GAM(\alpha, \alpha)$ with $\alpha = e^\eta$. A numerical procedure based on maximum likelihood is used to estimate the parameters of the model. The numerical procedure is iterative and uses a bisection root search method for estimating the gamma parameter η , least squares regression to estimate the Weibull ln-scale parameters β_0 through β_2 and an approximate Newton procedure to estimate the Weibull ln-shape parameters $\{\xi_i\}$. Simulation is used to study the behavior of the estimates for small sample sizes. Generally, the numerical estimation procedures work well. There are few numerical problems. All

the mean biases and mean square errors of the estimates tend to be small compared to the theoretical inputs to the simulation.

Topics for future research include replacing the one parameter gamma distribution with the two parameter gamma distribution in the hierarchical model and exploring the use of EM methodology to estimate model parameters. The two parameter gamma distribution arises as the posterior distribution of the current model. Another area is to study using the fitted hierarchical gamma/Weibull model to predict future performance.

It is hoped that the hierarchical gamma/Weibull regression model will be a useful tool to describe and predict one aspect of the effect of human performance on the battlefield.

LIST OF REFERENCES

1. Gaver, D. P., Jacobs, P. A., and O'Muircheartaigh, I., *Hierarchical Models for Target Detection Times*, Naval Postgraduate School Technical Report. To appear.
2. Bae, Deok Hwan, *Models for Target Detection Times*, Naval Postgraduate School Thesis, September 1989.
3. Johnson, L. W. and Riess, R. D., *Numerical Analysis*, Addison-Wesley, Inc. New York, 1977.
4. Peter A. Stark, *Introduction to Numerical methods*, Macmillan Publishing Co., Inc. New York, 1970.
5. Lewis, P. A. and Uribe, L., *The New Naval Postgraduate School Random Number Package LLRANDOM II*, Naval Postgraduate School Technical Report, NPS55-81-005, Monterey, CA, 1981.
6. Leonard Gilman and Allen J. Rose, *APL An Interactive Approach, Third Edition*, JOHN WILEY & SONS, Inc. New York, 1984.
7. P. McCullagh and J.A. Nelder, *Generalized Linear Models*, Chapman and Hall, New York, 1983.

APPENDIX A. STATISTICS AND GRAPHICS FOR THE ESTIMATE OF η IN THE SINGLE PARAMETER GAMMA MODEL

Table 1. Mean Bias, Mean Square Error, Std. Error and Uncensored Level for η at $O = 10$ in the Gamma Model

Number of Observers	Number of Targets	$\hat{\eta}$		UC
		M.B(S.E)	M.S.E(S.E)	
15	15	0.0727 (0.0497)	0.2477 (0.0693)	81%
	25	0.1027 (0.0436)	0.1967 (0.0269)	82%
	35	0.067 (0.0466)	0.2173 (0.0569)	82%
25	15	0.1003 (0.0311)	0.1051 (0.0167)	82%
	25	0.0566 (0.0304)	0.0935 (0.012)	82%
	35	0.0682 (0.0274)	0.0784 (0.0108)	82%
35	15	0.058 (0.0281)	0.0807 (0.0129)	82%
	25	0.0088 (0.0253)	0.0626 (0.0092)	82%
	35	0.0472 (0.0221)	0.05 (0.0103)	82%

Table 2. Mean Bias, Mean Square Error, Std. Error and Uncensored Level for η at $O = 15$ in the Gamma Model

Number of Observers	Number of Targets	$\hat{\eta}$		UC
		M.B(S.E)	M.S.E(S.E)	
15	15	0.0614 (0.0434)	0.1886 (0.0413)	89%
	25	0.0884 (0.0428)	0.1875 (0.026)	90%
	35	0.053 (0.0452)	0.2032 (0.052)	90%
25	15	0.094 (0.0305)	0.0998 (0.0151)	90%
	25	0.0517 (0.03)	0.0911 (0.0115)	90%
	35	0.0706 (0.0274)	0.0785 (0.0103)	90%
35	15	0.0519 (0.0277)	0.0781 (0.0132)	90%
	25	0.0085 (0.0246)	0.0593 (0.0086)	90%
	35	0.0471 (0.0216)	0.0481 (0.009)	90%

Table 3. Tendencies of Mean Bias and Mean Square Error for η at $O = 10$ in the Gamma Model

	M.B(M.S.E) of $\hat{\eta}$		
	15 TGT	25 TGT	35 TGT
15 OBS	0.073(0.248)	0.103(0.197)	0.067(0.217)
25 OBS	0.1 (0.105)	0.057(0.094)	0.068(0.078)
35 OBS	0.058(0.081)	0.009(0.063)	0.047(0.05)




Figure 1. The Tendencies of the Bias of η with changes in OBS and TGT at $O = 10$ in the Gamma Model: M = number of observers ; N = number of targets.




Figure 2. The Tendencies of the Bias of η with changes in OBS and TGT at $O = 10$ in the Gamma Model: M = number of observers ; N = number of targets.




Figure 3. The Tendencies of the Bias of η with changes in OBS and TGT at $O = 10$ in the Gamma Model: M = number of observers ; N = number of targets.




Figure 4. Comparison of the Bias of η between Different numbers of Observers with $TGT = 15$ at $O = 10$ in the Gamma Model: TGT = the number of targets.




Figure 5. Comparison of the Bias of η between Different numbers of Observers with TGT = 25 at O=10 in the Gamma Model: TGT = the number of targets.




Figure 6. Comparison of the Bias of η between Different numbers of Observers with $TGT = 35$ at $O = 10$ in the Gamma Model: TGT = the number of targets.

Table 4. Tendencies of Mean Bias and Mean Square Error for η at $O = 15$ in the Gamma Model

	M.B(M.S.E) of $\hat{\eta}$		
	15 TGT	25 TGT	35 TGT
15 OBS	0.061(0.189)	0.088(0.188)	0.053(0.203)
25 OBS	0.094(0.1)	0.052(0.091)	0.071(0.078)
35 OBS	0.052(0.078)	0.008(0.059)	0.047(0.048)




Figure 7. The Tendencies of the Bias of η with changes in OBS and TGT at $O = 15$ in the Gamma Model: M = number of observers ; N = number of targets.




Figure 8. The Tendencies of the Bias of η with changes in OBS and TGT at $O=15$ in the Gamma Model: M = number of observers ; N = number of targets.




Figure 9. The Tendencies of the Bias of η with changes in OBS and TGT at $O=15$ in the Gamma Model: M = number of observers ; N = number of targets.




Figure 10. Comparison of the Bias of η between Different numbers of Observers with TGT = 15 at O = 15 in the Gamma Model: TGT = the number of targets.




Figure 11. Comparison of the Bias of η between Different numbers of Observers with $TGT = 25$ at $O = 15$ in the Gamma Model: TGT = the number of targets.




Figure 12. Comparison of the Bias of η between Different numbers of Observers with $TGT = 35$ at $O = 15$ in the Gamma Model: TGT = the number of targets.

APPENDIX B. STATISTICS AND GRAPHICS FOR ALL THE ESTIMATES IN GAMMA/WEIBULL REGRESSION MODEL

1. TABLES AND HISTOGRAMS FOR η

Table 5. Mean Bias, Mean Square Error and Std. Error for η at $O = 10$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\eta}$	
		M.B(S.E)	M.S.E(S.E)
15	15	.1770(.0552)	.3299(.0892)
	25	.1573(.0455)	.2274(.0445)
	35	.0761(.0440)	.1955(.0290)
25	15	.0947(.0419)	.1811(.0420)
	25	.0571(.0315)	.1003(.0206)
	35	.1422(.0324)	.1228(.0208)
35	15	.1074(.0352)	.1328(.0216)
	25	.0658(.0307)	.0968(.0129)
	35	.0940(.0223)	.0577(.0070)

Table 6. Mean Bias, Mean Square Error and Std. Error for η at $O = 15$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\eta}$	
		M.B(S.E)	M.S.E(S.E)
15	15	.2107(.0561)	.3524(.1023)
	25	.1934(.0451)	.2365(.0458)
	35	.1117(.0442)	.2038(.0319)
25	15	.1170(.0418)	.1851(.0454)
	25	.0755(.0314)	.1021(.0222)
	35	.1434(.0314)	.1174(.0197)
35	15	.1450(.0336)	.1315(.0228)
	25	.0800(.0306)	.0982(.0133)
	35	.0968(.0219)	.0566(.0072)

Table 7. Tendencies of Mean Bias and Mean Square Error for η at $O = 10$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\eta}$		
	15 TGT	25 TGT	35 TGT
15 OBS	.177(.330)	.157(.227)	.076(.196)
25 OBS	.095(.181)	.057(.100)	.142(.123)
35 OBS	.107(.133)	.066(.097)	.094(.058)




Figure 13. Comparison of the Bias of η between Different numbers of Observers with TGT=15 at $O = 10$ in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 14. Comparison of the Bias of η between Different numbers of Observers with TGT=25 at O=10 in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 15. Comparison of the Bias of η between Different numbers of Observers with $TGT=35$ at $O=10$ in the GAM/WEI Regression Model: TGT = the number of targets.

Table 8. Tendencies of Mean Bias and Mean Square Error for η at $O = 15$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\eta}$		
	15 TGT	25 TGT	35 TGT
15 OBS.	.211(.352)	.193(.236)	.112(.204)
25 OBS.	.117(.185)	.075(.102)	.143(.117)
35 OBS.	.145(.132)	.080(.098)	.097(.057)




Figure 16. Comparison of the Bias of η between Different numbers of Observers with TGT=15 at $O=15$ in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 17. Comparison of the Bias of η between Different numbers of Observers with TGT=25 at O=15 in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 18. Comparison of the Bias of η between Different numbers of Observers with TGT=35 at O=15 in the GAM/WEI Regression Model: TGT = the number of targets.

2. TABLES AND HISTOGRAMS FOR β_0

Table 9. Mean Bias, Mean Square Error and Std. Error for β_0 at $O = 10$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\beta}_0$	
		M.B(S.E)	M.S.E(S.E)
15	15	-.0605(.0317)	.1021(.0123)
	25	-.0234(.0278)	.0764(.0109)
	35	.0304(.0223)	.0499(.0067)
25	15	.0458(.0238)	.0575(.0081)
	25	-.0167(.0222)	.0487(.0052)
	35	-.0094(.0197)	.0383(.0056)
35	15	-.0067(.0198)	.0386(.0067)
	25	-.0398(.0176)	.0319(.0047)
	35	.0062(.0152)	.0227(.0027)

Table 10. Mean Bias, Mean Square Error and Std. Error for β_0 at $O = 15$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\beta}_0$	
		M.B(S.E)	M.S.E(S.E)
15	15	-.0609(.0316)	.1015(.0127)
	25	-.0234(.0271)	.0725(.0095)
	35	.0263(.0224)	.0498(.0063)
25	15	.0365(.0244)	.0595(.0073)
	25	-.0292(.0210)	.0439(.0049)
	35	-.0185(.0184)	.0335(.0053)
35	15	-.0043(.0191)	.0358(.0066)
	25	-.0454(.0176)	.0324(.0041)
	35	-.0021(.0138)	.0187(.0023)

Table 11. Tendencies of Mean Bias and Mean Square Error for β_0 at $O = 10$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\beta}_0$		
	15 TGT	25 TGT	35 TGT
15 OBS	-.061(.102)	-.023(.076)	.030(.050)
25 OBS	.046(.051)	-.017(.049)	-.009(.038)
35 OBS	-.007(.039)	-.040(.032)	.006(.023)




Figure 19. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 15 at $O = 10$ in the GAM/WEI Regression Model: OBS = the number of observers




Figure 20. Comparison of the Bias of β_0 between Different numbers of targets with OBS=25 at O=10 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 21. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 35 at O = 10 in the GAM/WEI Regression Model: OBS = the number of observers

Table 12. Tendencies of Mean Bias and Mean Square Error for β_0 at $O = 15$ in the GAM/WEI Regression Model.

	M.B(M.S.E) of $\hat{\beta}_0$		
	15 TGT	25 TGT	35 TGT
15 OBS	-.061(.102)	-.023(.073)	.026(.050)
25 OBS	.037(.060)	-.029(.044)	-.018(.033)
35 OBS	-.004(.036)	-.045(.032)	-.002(.019)




Figure 22. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 15 at O = 15 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 23. Comparison of the Bias of β_0 between Different numbers of targets with OBS=25 at O=15 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 24. Comparison of the Bias of β_0 between Different numbers of targets with OBS = 35 at O = 15 in the GAM/WEI Regression Model: OBS = the number of observers

3. TABLES AND HISTOGRAMS FOR β_1

Table 13. Mean Bias, Mean Square Error and Std. Error for β_1 at $O = 10$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\beta}_1$	
		M.B(S.E)	M.S.E(S.E)
15	15	.0180(.0157)	.0243(.0032)
	25	-.0002(.0140)	.0193(.0026)
	35	-.0010(.0112)	.0122(.0017)
25	15	-.0154(.0132)	.0173(.0026)
	25	-.0006(.0096)	.0091(.0014)
	35	-.0036(.0080)	.0063(.0010)
35	15	.0188(.0126)	.0160(.0020)
	25	.0094(.0096)	.0091(.0011)
	35	.0006(.0069)	.0047(.0006)

Table 14. Mean Bias, Mean Square Error and Std. Error for β_1 at $O = 15$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\beta}_1$	
		M.B(S.E)	M.S.E(S.E)
15	15	.0079(.0145)	.0205(.0028)
	25	-.0060(.0127)	.0158(.0022)
	35	.0076(.0107)	.0113(.0014)
25	15	-.0214(.0127)	.0162(.0024)
	25	.0036(.0090)	.0079(.0013)
	35	-.0046(.0075)	.0055(.0009)
35	15	.0150(.0121)	.0146(.0018)
	25	.0077(.0092)	.0084(.0010)
	35	.0009(.0067)	.0043(.0005)

Table 15. Tendencies of Mean Bias and Mean Square Error for β_1 at $O = 10$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\beta}_1$		
	15 TGT	25 TGT	35 TGT
15 OBS	.018(.024)	.000(.019)	-.001(.012)
25 OBS	-.015(.017)	-.001(.009)	-.004(.006)
35 OBS	.019(.016)	.009(.009)	.001(.005)




Figure 25. Comparison of the Bias of β_1 between Different numbers of targets with OBS=15 at O=10 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 26. Comparison of the Bias of β_1 between Different numbers of targets with OBS=25 at O=10 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 27. Comparison of the Bias of β_1 between Different numbers of targets with OBS = 35 at O = 10 in the GAM/WEI Regression Model: OBS = the number of observers

Table 16. Tendencies of Mean Bias and Mean Square Error for β_1 at $O = 15$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\beta}_1$		
	15 TGT	25 TGT	35 TGT
15 OBS	.008(.021)	-.006(.016)	.008(.011)
25 OBS	-.021(.016)	.004(.008)	-.005(.006)
35 OBS	.015(.015)	.008(.008)	.001(.004)




Figure 28. Comparison of the Bias of β_1 between Different numbers of targets with OBS=15 at O=15 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 29. Comparison of the Bias of β_1 between Different numbers of targets with OBS=25 at O=15 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 30. Comparison of the Bias of β_1 between Different numbers of targets with OBS=35 at O=15 in the GAM/WEI Regression Model: OBS = the number of observers

4. TABLES AND HISTOGRAMS FOR β_2

Table 17. Mean Bias, Mean Square Error and Std. Error for β_2 at $O = 10$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\beta}_2$	
		M.B(S.E)	M.S.E(S.E)
15	15	.0167(.0097)	.0096(.0012)
	25	.0057(.0075)	.0056(.0008)
	35	-.0097(.0053)	.0029(.0004)
25	15	-.0051(.0073)	.0053(.0008)
	25	.0040(.0054)	.0029(.0004)
	35	.0020(.0046)	.0021(.0003)
35	15	.0002(.0062)	.0038(.0006)
	25	.0083(.0044)	.0020(.0003)
	35	-.0046(.0033)	.0011(.0001)

Table 18. Mean Bias, Mean Square Error and Std. Error for β_2 at $O = 15$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\beta}_2$	
		M.B(S.E)	M.S.E(S.E)
15	15	.0185(.0091)	.0084(.0012)
	25	.0055(.0070)	.0049(.0006)
	35	-.0124(.0052)	.0028(.0004)
25	15	-.0015(.0069)	.0046(.0006)
	25	.0072(.0050)	.0025(.0003)
	35	.0024(.0042)	.0018(.0002)
35	15	-.0016(.0057)	.0032(.0005)
	25	.0086(.0045)	.0020(.0003)
	35	-.0007(.0032)	.0010(.0001)

Table 19. Tendencies of Mean Bias and Mean Square Error for β_2 at $O = 10$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\beta}_2$		
	15 TGT	25 TGT	35 TGT
15 OBS	.017(.010)	.006(.006)	-.010(.003)
25 OBS	-.005(.005)	.004(.003)	.002(.002)
35 OBS	.000(.004)	.008(.002)	-.005(.001)




Figure 31. Comparison of the Bias of β_2 between Different numbers of targets with OBS=15 at $O=10$ in the GAM/WEI Regression Model: OBS = the number of observers




Figure 32. Comparison of the Bias of β_2 between Different numbers of targets with OBS=25 at O=10 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 33. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 35 at O = 10 in the GAM/WEI Regression Model: OBS = the number of observers

Table 20. Tendencies of Mean Bias and Mean Square Error for β_2 at O = 15 in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\beta}_2$		
	15 TGT	25 TGT	35 TGT
15 OBS	.018(.008)	.006(.005)	-.012(.003)
25 OBS	-.001(.005)	.007(.002)	.002(.002)
35 OBS	-.002(.003)	.009(.002)	-.001(.001)




Figure 34. Comparison of the Bias of β_2 between Different numbers of targets with OBS = 15 at O = 15 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 35. Comparison of the Bias of β_2 between Different numbers of targets with OBS=25 at O=15 in the GAM/WEI Regression Model: OBS = the number of observers




Figure 36. Comparison of the Bias of β_2 between Different numbers of targets with OBS=35 at O=15 in the GAM/WEI Regression Model: OBS = the number of observers

5. TABLES AND HISTOGRAMS FOR ξ

Table 21. Mean Bias, Mean Square Error and Std. Error for ξ at $O = 10$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\xi}$	
		M.B(S.E)	M.S.E(S.E)
15	15	-.0318(.0067)	.0054(.0007)
	25	-.0158(.0056)	.0033(.0005)
	35	-.0131(.0048)	.0024(.0003)
25	15	-.0226(.0053)	.0033(.0005)
	25	-.0078(.0036)	.0013(.0002)
	35	-.0025(.0038)	.0014(.0002)
35	15	-.0196(.0044)	.0023(.0003)
	25	-.0065(.0036)	.0013(.0002)
	35	-.0071(.0028)	.0008(.0001)

Table 22. Mean Bias, Mean Square Error and Std. Error for ξ at $O = 15$ in the GAM/WEI Regression Model

Number of Observers	Number of Targets	$\hat{\xi}$	
		M.B(S.E)	M.S.E(S.E)
15	15	-.0349(.0064)	.0052(.0011)
	25	-.0147(.0051)	.0027(.0005)
	35	-.0129(.0041)	.0018(.0002)
25	15	-.0181(.0047)	.0025(.0003)
	25	-.0089(.0037)	.0014(.0002)
	35	-.0027(.0035)	.0012(.0002)
35	15	-.0192(.0041)	.0020(.0003)
	25	-.0049(.0033)	.0011(.0002)
	35	-.0083(.0025)	.0007(.0001)

Table 23. Tendencies of Mean Bias and Mean Square Error for ξ at $O = 10$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\xi}$		
	15 TGT	25 TGT	35 TGT
15 OBS	-.032(.005)	-.016(.003)	-.013(.002)
25 OBS	-.023(.003)	-.008(.001)	-.003(.001)
35 OBS	-.020(.002)	-.007(.001)	-.007(.001)




Figure 37. Comparison of the Bias of ξ between Different numbers of Observers with TGT=15 at $O=10$ in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 38. Comparison of the Bias of ξ between Different numbers of Observers with TGT=25 at O=10 in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 39. Comparison of the Bias of ξ between Different numbers of Observers with TGT=35 at O=10 in the GAM/WEI Regression Model: TGT = the number of targets.

Table 24. Tendencies of Mean-Bias and Mean Square Error for ξ at $O = 15$ in the GAM/WEI Regression Model

	M.B(M.S.E) of $\hat{\xi}$		
	15 TGT	25 TGT	35 TGT
15 OBS	-.035(.005)	-.015(.003)	-.013(.002)
25 OBS	-.018(.002)	-.009(.001)	-.003(.001)
35 OBS	-.019(.002)	-.005(.001)	-.008(.001)




Figure 40. Comparison of the Bias of ξ between Different numbers of Observers with TGT=15 at $O=15$ in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 41. Comparison of the Bias of ξ between Different numbers of Observers with $TGT=25$ at $O=15$ in the GAM/WEI Regression Model: TGT = the number of targets.




Figure 42. Comparison of the Bias of ξ between Different numbers of Observers with $TGT=35$ at $O=15$ in the GAM/WEI Regression Model: TGT = the number of targets.

APPENDIX C. SIMULATION PROGRAM FOR THE ESTIMATE OF η IN THE SINGLE PARAMETER GAMMA MODEL

```

V R←SIMULA1;M;N;O;I;J;K;L;ETA;OUT;PAGE;AVUCL;MB;SEMB;MSE;SEMSE
;RRR
[1]  AAAAAAAAA KEY GLOBAL VARIABLES : BOOK1 ; BOOK2 ; UCL AAAAAAAAA
[2]  AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA DATA INPUT AAAAAAAAAAAAAAAAA
[3]  M← 15 25 35
[4]  N← 15 25 35
[5]  O← 10 15
[6]  ETA←1
[7]  I←J←K←L←0
[8]  AAAAAAAAAAAAAAAAA START LOOPING THE SIMULATION 1 AAAAAAAAA
[9]  LOOPI:I←I+1
[10] LOOPJ:J←J+1
[11] LOOPK:K←K+1
[12] AAAAAAAAAAAAAAAAA RESET THE RANDOM NUMBER SEED AAAAAAAAA
[13] R←O[I],M[J],N[K]
[14] RRL←466801743
[15] LOOPL:L←L+1
[16] AAAAAAAAAAAAAAAAA CALL GAMMA ESTIMATE FUNCTION AAAAAAAAA
[17] RRR←RRL
[18] OUT←GAMMA(O[I],M[J],N[K])
[19] AAAAAAA RECORD UCL AND ALL THE ESTIMATES IN REPLICATIONS AAAAA
[20] →((L=1),L>1)/INTZ1,GO1
[21] INTZ1:PAGE←(1,ρOUT)ρ(OUT-0,ETA)
[22] →LOOPL
[23] GO1:PAGE←PAGE,[1](OUT-0,ETA)
[24] TESTL:
[25] →(L<100)/LOOPL
[26] AAAAAAA COMPUTE AVE. UNCENSORED LEVEL AND ALL THE STATISTICS AA
[27] AVUCL←((+/PAGE[;1])÷L)
[28] MB←((+/PAGE[;2])÷L)
[29] SEMB←((+/((PAGE[;2]-MB)*2)÷(L×L-2))*0.5)
[30] MSE←((+/PAGE[;2]*2)÷L)
[31] SEMSE←((+/((PAGE[;2]*2)-MSE)*2)÷(L×L-2))*0.5
[32] L←0
[33] AAAAAAA KEEP ALL THE INFORMATIONS FOR GRAPHICS AND TABLES AA
[34] →(((I×J×K)=1),(I×J×K)≠1)/INTZ2,GO2
[35] INTZ2:
[36] BOOK1←(1,ρPAGE[;2])ρPAGE[;2]
[37] BOOK2←(1,4)ρ(MB,SEMB,MSE,SEMSE)
[38] UCL←AVUCL
[39] →TESTK
[40] GO2:
[41] BOOK1←BOOK1,[1] PAGE[;2]
[42] BOOK2←BOOK2,[1](MB,SEMB,MSE,SEMSE)

```

[43] $UCL \leftarrow UCL, AVUCL$
[44] $TESTK:$
[45] $\rightarrow (K < pN) / LOOPK$
[46] $K \leftarrow 0$
[47] $\rightarrow (J < pM) / LOOPJ$
[48] $J \leftarrow 0$
[49] $\rightarrow (I < pO) / LOOPI$
 ∇

```

      V R←GAMMA INPUT;M;N;O;ETA;ETA0;XI;XIO;A;WW;MU;U;Y;DELTA;C;XB;RR
      ;M1;M2;T;DATA;K;FL;FM;FR;LB;MP;RB;S
[1]  AAAAAAAAAA SIMULATION INPUTS AND THEORETICAL PARAMETERS AAAAA
[2]  M←INPUT[2]
[3]  N←INPUT[3]
[4]  O←(M,N)ρINPUT[1]
[5]  ETA←1
[6]  MU←4.2
[7]  XIO←Q(N,M)ρXI←MpO
[8]  AAAAAAAAAAAAAAAAAA RANDOM NUMBER GENERATIONS AAAAAAAAAAAAAAAAAA
[9]  A←Q(N,M)ρM GAMRAND((∗ETA),+∗ETA)
[10] WW←(M,N)ρ(M×N) EXPRAND 1
[11] AAAAAAAAAAAAAAAAAA DETECTION TIME AND UNCENSORING DATA AAAAAAAAAAAAAAAAAA
[12] U←MU×(WW+A)×(∗XIO)
[13] A←WW+10
[14] Y←(⊗U)⊔(⊗O)
[15] C←+/[2] DELTA←(⊗U)≤(⊗O)
[16] O←10
[17] XB←⊗MU
[18] AAAAAAAAAAAAAAAAAA INITIAL CONDITION FOR GAMMA PARAMETER ,ETA AAAAAA
[19] RR←(Y-XB)×∗(1×XIO)
[20] M1←0.5772
[21] M2←(+ / + / DELTA × RR × 2) + ( + / C)
[22] ETA0←0
[23] + ( O > T ← ( ( ( o 1 ) × 2 ) + 6 ) + ( M 2 × 2 ) - ( M 1 × 2 ) ) / B I S E C T I O N
[24] ETA0←1×⊗T
[25] AAAAA FIND BOUNDS AND BISECTION SEARCH FOR GAMMA PARAMETER,ETA AA
[26] BISECTION:K←0
[27] DATA←(S←+/[2]×RR),C
[28] FM←ETA0 FVALUE DATA
[29] AAAAAAAAAAAAAAAAAA FIND THE RIGHT AND LEFT BOUND FOR ETA AAAAAAAAAAAAAAAAAA
[30] BOUND:K←K+1
[31] FL←(LB←ETA0-K×0.5) FVALUE DATA
[32] FR←(RB←ETA0+K×0.5) FVALUE DATA
[33] + ( ( ( × F M ) × ( × F L ) ) < 0 ) / S E T R B
[34] + ( ( ( × F M ) × ( × F R ) ) < 0 ) / S E T L B
[35] →BOUND
[36] SETRB:RB←ETA0
[37] →BISECT
[38] SETLB:LB←ETA0
[39] AAAAAAAAAAAAAAAAAA PERFORM THE BISECTION SEARCH FOR ETA AAAAAAAAAAAAAAAAAA
[40] BISECT:
[41] FM←(MP←(LB+RB)+2) FVALUE DATA
[42] FL←LB FVALUE DATA
[43] + ( ( R B - L B ) < ( | 0 . 0 1 × M P ) ) / E N D
[44] + ( ( ( ( × F M ) × ( × F L ) ) < 0 ) , ( ( ( × F M ) × ( × F L ) ) > 0 ) ) / S W A P R , S W A P L
[45] SWAPR:RB←MP
[46] →BISECT
[47] SWAPL:LB←MP
[48] →BISECT

```

```

[49]  AAAAAAAAAAAAAAAAAAAAAAAAAA  OUTPUT THE ESTIMATES  AAAAAAAAAAAAAAAAAAAAA
[50]  END:
[51]  ETA0←(RB+LB)÷2
[52]  R←((+/C)÷M×N),ETA0
      ∇

```

```

      ∇ R←ETA FVALUE DATA;M;S;C;SUBF;I
[1]  M←(ρDATA)÷2
[2]  S←M÷DATA
[3]  C←M÷DATA
[4]  SUBF←0
[5]  I←0
[6]  LOOP:I←I+1
[7]  →(C[I]=0)/CHECKI
[8]  SUBF←SUBF+(+/÷(*ETA)+(1C[I])-1)
[9]  CHECKI:
[10] →(I<M)/LOOP
[11] R←(*ETA)×SUBF+(+/ETA+((S-C)÷S+*ETA)+-1×⊗(S+*ETA))
      ∇

```

APPENDIX D. SIMULATION PROGRAM FOR THE ESTIMATES OF ALL PARAMETERS IN GAMMA/WEIBULL REGRESSION MODEL

```

V R+SIMULA2;M;N;O;I;J;K;L;X;E;ETA;B0;B1;B2;XI;OUT;PAGE;AVUCEX;MB
;SEMB;MSE;SEMSE;RRR
[1] AAAAAAAAAA KEY GLOBAL VARIABLES : BOOK1 ; BOOK2 ; UCLEX AAAAAAAAAA
[2] AAAAAAAAAA DATA INPUT AAAAAAAAAA
[3] M+ 15 25 35
[4] N+ 15 25 35
[5] O+ 10 15
[6] ETA+1
[7] B0+0.8
[8] B1+0.2
[9] B2+0.5
[10] XI+0
[11] AAAAAAAAAA START LOOPING THE SIMULATION AAAAAAAAAA
[12] I+J+K+L+E+X+0
[13] LOOPI:I+I+1
[14] LOOPJ:J+J+1
[15] LOOPK:K+K+1
[16] AAAAAAAAAA RESET THE RANDOM NUMBER SEED AAAAAAAAAA
[17] R+O[I],M[J],N[K]
[18] RRL+466801743
[19] LOOPL:L+L+1
[20] AAAAAAAAAA CALL GAMMA/WEIBULL REGRESSION FUNCTION AAAAAAAAAA
[21] RRR+RRL
[22] OUT+GAMWEI(O[I],M[J],N[K])
[23] AAAAAAAAAA SKIP THE INSUFFICIENT VARIABILITY CASE FOR GAMMA AA
[24] +(OUT[2]*~99999)/CHECKETA
[25] E+E+1
[26] L+L-1
[27] +TESTL
[28] AAAAAA RECORD THE RANDOM SEED FOR THE ILL CONDITION OF ETA AAA
[29] CHECKETA:
[30] +(OUT[2]*~100000)/GOGO
[31] R+'BAD ETA'
[32] R+RRR
[33] L+L-1
[34] +TESTL
[35] AAAAAA RECORD UCL AND ALL THE ESTIMATES IN REPLICATIONS AAA
[36] GOGO:
[37] +((L=1),L>1)/INTZ1,G01
[38] INTZ1:PAGE+(1,(pOUT)-1)p((-1+OUT)-0,ETA,B0,B1,B2,XI)
[39] X+X+((-1+OUT)
[40] +LOOPL
[41] G01:PAGE+PAGE,[1]((-1+OUT)-0,ETA,B0,B1,B2,XI)
[42] X+X+((-1+OUT)

```

```

[43] AAAAAAAAA RECORD THE RANDOM SEED FOR THE ILL CASE OF XI AAAAAAAAA
[44] →((-1+OUT)=0)/TESTL
[45] □←'BAD XI'
[46] □←RRR
[47] AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
[48] TESTL:
[49] →(L<100)/LOOPL
[50] AAAAAAA COMPUTE AVE. UNCENSORED LEVEL AND ALL THE STATISTICS A
[51] AVUCEX←((+/PAGE[;1])÷L),E,X
[52] MB←(+/[1] PAGE← 0 1 ↓PAGE)÷L
[53] SEMB←((+/[1] (PAGE-(pPAGE)pMB)*2)+(L×L-2))*0.5
[54] MSE←(+/[1] PAGE*2)÷L
[55] SEMSE←((+/[1] ((PAGE*2)-(pPAGE)pMSE)*2)+(L×L-2))*0.5
[56] L←E+X←0
[57] AAAAAAA KEEP ALL THE INFORMATIONS FOR GRAPHICS AND TABLES AAA
[58] →(((I×J×K)=1),(I×J×K)≠1)/INTZ2,GO2
[59] INTZ2:
[60] BOOK1←(1,pPAGE)pPAGE
[61] BOOK2←(1,4,pMB)p(MB,SEMB,MSE,SEMSE)
[62] UCLEX← 1 3 pAVUCEX
[63] →TESTK
[64] GO2:
[65] BOOK1+BOOK1,[1] PAGE
[66] BOOK2+BOOK2,[1] ((4,pMB)pMB,SEMB,MSE,SEMSE)
[67] UCLEX+UCLEX,[1] AVUCEX
[68] TESTK:
[69] →(K<pN)/LOOPK
[70] K←0
[71] →(J<pM)/LOOPJ
[72] J←0
[73] →(I<pO)/LOOPI
V

```

```

VGAMWEI[ ]V
V R←GAMWEI INPUT;M;N;O;ETA;ETA0;B;B0;B1;B2;XI;XIO;A;WW;X1;X2
;MU;U;Y;DELTA;C;XB;RR;W;U0;U1;U2;UU;Z;I;VAR;M1;M2;T;DATA;K
;FL;FLO;FM;FR;FRO;LB;MP;RB;CS;H;DH;DIFFXI;BK;S;J;X;XX;BADXI
[1]  AAAAAAAAAA SIMULATION INPUTS AND THEORETICAL PARAMETERS AAAAAA
[2]  M←INPUT[2]
[3]  N←INPUT[3]
[4]  O←(M,N)ρINPUT[1]
[5]  ETA←1
[6]  B0←0.8
[7]  B1←0.2
[8]  B2←0.5
[9]  XIO←q(N,M)ρXI←Mp0
[10] AAAAAAAAAA RANDOM NUMBER GENERATIONS AAAAAAAAAA
[11] A←q(N,M)ρM GAMRAND(*ETA),+*ETA)
[12] WW←(M,N)ρ(M×N) EXPRAND 1
[13] X1←(M,N)ρ(M×N) NORRAND 1 0.5
[14] X2←(M,N)ρ(M×N) NORRAND 2 1
[15] AAAAAAAAAA DETECTION TIME AND UNCENSORING DATA AAAAAAAAAA
[16] MU←*(B0+(B1×X1)+B2×X2)
[17] U←MU×(WW+A)*(XIO)
[18] A×WW←MU←10
[19] Y←(⊗U)⊗(⊗0)
[20] C←+/[2] DELTA←(⊗U)≤(⊗0)
[21] O←10
[22] AAAAAAAAAA INITIAL ESTIMATES WITHOUT GAMMA R.V. INVOLVED AAAAAA
[23] XB←⊗U
[24] I←0
[25] INTZ:I←I+1
[26] W←*(Y-XB)**(-1×XIO))*0.5
[27] U0←((M,N)ρ1)×W*(-1×XIO)
[28] U1←X1×W*(-1×XIO)
[29] U2←X2×W*(-1×XIO)
[30] UU←q(3,M×N)ρ(,U0),(,U1),(,U2)
[31] U0←U1←U2←10
[32] Z←q,(W×XB*(-1×XIO))+((-1×DELTA)+W*2)÷W
[33] W←10
[34] B←Z⊗UU
[35] Z←UU←10
[36] BK←B
[37] XB←B[1]+(B[2]×X1)+B[3]×X2
[38] RR←(Y-XB)**(-1×XIO)
[39] XIO←q(N,M)ρXI←XI+DIFFXI←(C-+/RR×((-1×DELTA)+*RR))÷
((-1×C)-+/((RR*2)*RR)
[40] →(I<2)/INTZ
[41] AAAAAAAAAA VARIABILITY CHECK FOR UNCENSORED TIMES AAAAAAAAAA
[42] VAR←(+/+/(-1+ (U÷(*XB))*(-1×XIO))*2)÷(-1+M×N)
[43] U←10
[44] ETA0←-99999
[45] X←Mp1

```



```

[46] J←0
[47] →(VAR<1)/END
[48] AAAAAAAAAAAAA INITIAL CONDITION FOR GAMMA PARAMETER ,ETA AAAAAA
[49] RR←(Y-XB)**(-1×XIO)
[50] M1←-0.5772
[51] M2←(+ / + / DELTA×RR*2) ÷ (+ / C)
[52] ETA0←0
[53] →(0>T÷(((01)*2)÷-6)+(M2*2)-(M1*2))/READY
[54] ETA0←-1×T
[55] READY:BADXI←0
[56] AA LARGE RECURSIVE LOOP INCLUDING ESTIMATE OF GAMMA PARAMETER,ETA
[57] RECURSIVE:K←0
[58] J←J+1
[59] DATA←(S÷+[2]*RR),C
[60] FM←ETA0 FVALUE DATA
[61] AAAAAA BISECTION SEARCH METHOD APPLIED TO ESTIMATE ETA AAAAAA
[62] BOUND:K←K+1
[63] FL←(LB←ETA0-K×0.5) FVALUE DATA
[64] FR←(RB←ETA0+K×0.5) FVALUE DATA
[65] AAAAAAAAAAAAAAAAA DETECT ILL CONDITION FOR ETA AAAAAAAAAAAAAAAAA
[66] →(K≤10)/CHECKLRB
[67] →(∼(((|FRO-FR)÷0.5)<0.01)^(|FR|<0.1))/CHECKLRB
[68] ETA0←-100000
[69] →END
[70] AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
[71] CHECKLRB:
[72] →(((×FM)×(×FL))<0)/SETRB
[73] →(((×FM)×(×FR))<0)/SETLB
[74] FLO←FL
[75] FRO←FR
[76] →BOUND
[77] SETRB:RB←ETA0
[78] →BISECT
[79] SETLB:LB←ETA0
[80] BISECT:
[81] FM←(MP←(LB+RB)÷2) FVALUE DATA
[82] FL←LB FVALUE DATA
[83] →((RB-LB)<(|0.01×MP))/REGRESS
[84] →((((×FM)×(×FL))<0),(((×FM)×(×FL))>0))/SWAPR,SWAPL
[85] SWAPR:RB←MP
[86] →BISECT
[87] SWAPL:LB←MP
[88] →BISECT
[89] AAAAA REGRESSION PROCEDURE FOR WEIBULL SCALE PARAMETERS,BETA'S AA
[90] REGRESS:
[91] ETA0←(RB+LB)÷2
[92] W←((CS÷Q(N,M)ρ((C+×ETA0)÷S+×ETA0))××RR)*0.5
[93] U0←((M,N)ρ1)×W×*(-1×XIO)
[94] U1←X1×W×*(-1×XIO)
[95] U2←X2×W×*(-1×XIO)
[96] UU←Q(3,M×N)ρ(,U0),(,U1),(,U2)

```

```

[97]  U0←U1←U2←10
[98]  Z←Q,(W×XB×*(-1×XIO))+((-1×DELTA)+W*2)÷W
[99]  B←ZUU
[100]  UU←Z←10
[101]  AAAAAAAAA NEWTON PROCEDURE FOR WEIBULL SHAPE PARAMETER,XI'S AAA
[102]  XB←B[1]+(B[2]×X1)+B[3]×X2
[103]  RR←(Y-XB)×*(-1×XIO)
[104]  H←(-1×C)+(+/[2] RR×((-1×DELTA)+CS×RR))
[105]  DH←(-1×C)-(+/[2] CS×(*RR)×RR*2)
[106]  AAAAAAAAAAAAA DETECT THE ILL CONDITION OF XI'S AAAAAAAAAAAAA
[107]  XX←~((|H)<0.1)^(|DH)<0.01)
[108]  →(~((×/XX)=0)∧BADXI=0)/NEWTONXI
[109]  BADXI←1
[110]  X←XX
[111]  NEWTONXI:
[112]  XIO←Q(N,M)ρXI←XI+DIFFXI←XX×(-1×H)+DH
[113]  AA TEST FOR STOPPING CRITERIA AND UPDATE THE VALUE OF VARIABLES
[114]  →((×/(|DIFFXI+XI),(BK-B)÷B)<0.01)∨(J≥50))/END
[115]  BK←B
[116]  RR←(Y-XB)×*(-1×XIO)
[117]  →RECURSIVE
[118]  AAAAAAAAA OUTPUT UNCENSORED LEVEL AND ALL ESTIMATES AAAAAAAAA
[119]  END:
[120]  R←((+/C)÷M×N),ETA0,B[1],B[2],B[3],((+/XI×X)÷(+/X)),+/~X
V

```

INITIAL DISTRIBUTION LIST

		No. Copies
1.	Defense Technical Information Center Cameron Station Alexandria, Virginia 22304-6145	2
2.	Library, Code 52 Naval Postgraduate School Monterey, California 93943-5002	2
3.	Professor P. A. Jacobs, Code ORJC Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	5
4.	Professor D. P. Gaver, Code ORGV Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	5
5.	Professor Samuel Parry, Code ORPY Department of Operations Research Naval Postgraduate School Monterey, California 93943-5000	1
6.	Lt. Col. Bard Mansager TRADOC Monterey, California 93945	1
7.	Epaminondas Hatzopoulos K. Palaiologou 5 Nea Smirni 17121 Athens, Greece	1
8.	Lt. Rogerio Guerra da Silveira Rua Lopes Quintas, 390, apt. 402 Jardim Botânico, Rio de Janeiro RJ 22240, Brazil	1

9. Wang Chia-Fu
No. 9, Aly. 30, Ln. 185, Central Rd. Sec. 4
Tu-Cheng 23604, Taipei County
Taiwan R.O.C 2
10. Won Wen-I
SMC 1830, NPS
Monterey, CA 93943 1